

Set Items Description

Ref	Items	RT	Index-term
E1	1		TETANUCLEATE
E2	1		TETANUKSEN
E3	13902	3	*TETANUS
E4	71		TETANUS --BLOOD --BL
E5	5		TETANUS --CEREBROSPINAL FLUID --CF
E6	30		TETANUS --CHEMICALLY INDUCED --CI
E7	12		TETANUS --CLASSIFICATION --CL
E8	393		TETANUS --COMPLICATIONS --CO
E9	21		TETANUS --CONGENITAL --CN
E10	338		TETANUS --DIAGNOSIS --DI
E11	3		TETANUS --DIET THERAPY --DH
E12	523		TETANUS --DRUG THERAPY --DT

Ref	Items	Type	RT	Index-term
R1	13902	3		*TETANUS
R2	4515	X		DC=C1.252.410.90.217.864. (TETANUS)
R3	904	R	6	TRISMUS
R4	2533	B	7	CLOSTRIDIUM INFECTIONS

Ref	Items	Type	RT	Index-term
R1	2533	7		*CLOSTRIDIUM INFECTIONS
R2	2533	X		DC=C1.252.410.90.217. (CLOSTRIDIUM INFECTIONS)
R3	122	B	9	BACILLACEAE INFECTIONS
R4	1785	N	10	BOTULISM
R5	3997	N	10	ENTEROCOLITIS, PSEUDOMEMBRANOUS
R6	279	N	3	ENTEROTOXEMIA
R7	1368	N	2	GAS GANGRENE
R8	13902	N	3	TETANUS

S1 13902 "TETANUS"
S2 2533 "CLOSTRIDIUM INFECTIONS"
S3 90754 THYROID
S4 38 S1 AND S3
S5 2 S2 AND S3

Ref	Items	RT	Index-term
E1	753		THYROTROPIC
E2	2		THYROTROPICAL
E3	25841	8	*THYROTROPIN
E4	167		THYROTROPIN --ADMINISTRATION AND DOSAGE --AD
E5	30		THYROTROPIN --ADVERSE EFFECTS --AE
E6	17		THYROTROPIN --ANALOGS AND DERIVATIVES --AA
E7	896		THYROTROPIN --ANALYSIS --AN
E8	255		THYROTROPIN --ANTAGONISTS AND INHIBITORS --AI
E9	288		THYROTROPIN --BIOSYNTHESIS --BI
E10	10117		THYROTROPIN --BLOOD --BL
E11	12		THYROTROPIN --CEREBROSPINAL FLUID --CF
E12	6		THYROTROPIN --CHEMICAL SYNTHESIS --CS

S6 25841 "THYROTROPIN"
S7 19 S6 AND (S1 OR S2)
S8 30150 "THYROID GLAND"

Ref	Items	RT	Index-term
E1	96		THYROID FUNCTION TESTS --VETERINARY --VE
E2	0	1	THYROID GALACTOSYLTRANSFERASE
E3	30150	5	*THYROID GLAND

Ref	Items	RT	Index-term
E4	595		THYROID GLAND --ABNORMALITIES --AB
E5	1135		THYROID GLAND --ANALYSIS --AN
E6	992		THYROID GLAND --ANATOMY AND HISTOLOGY --AH
E7	672		THYROID GLAND --BLOOD SUPPLY --BS
E8	369		THYROID GLAND --CHEMISTRY --CH
E9	2573		THYROID GLAND --CYTOLOGY --CY
E10	1		THYROID GLAND --DIAGNOSIS --DI
E11	5066		THYROID GLAND --DRUG EFFECTS --DE
E12	601		THYROID GLAND --EMBRYOLOGY --EM

Ref	Items	Type	RT	Index-term
R1	30150	5		*THYROID GLAND
R2	30150	X		DC=A6.407.900. (THYROID GLAND)
R3	221	R	3	STRUMA OVARII
R4	11014	R	3	THYROIDECTOMY
R5	217	R	3	ULTIMOBRANCHIAL BODY
R6	5409	B	30	ENDOCRINE GLANDS

S9 17616 CLOSTRID?
S10 7 S8 AND S9
S11 133 PHOSPHOLIPASE AND NEUROTOXIN
S12 27833 THYROX?
S13 2 S9 AND S12
S14 6426 BOTULI?
S15 0 S14 AND S12
S16 0 S14 AND S8
S17 18 S3 AND S14

4/6/1 11245935 21195143 PMID: 11298123
Intradermal skin test with diabetes specific antigens in patients with type 1 diabetes. Mar 2001

4/6/2 10726207 20373526 PMID: 10916814
[Preventive activities in primary health care: identifying the agreement among evidence-based guidelines] Actividades preventivas en atencion primaria: identificacion de areas de concordancia entre guias de practica clinica basadas en la evidencia. 2000

4/6/3 10722524 20424529 PMID: 10970112
Muscle reinnervation in hypothyroid rats. 1996

4/6/4 10042358 99134381 PMID: 9933650
Detection of novel carbohydrate binding activity of interleukin-1. Feb 12 1999

4/6/5 09732112 98204670 PMID: 9545109
Induction of oral tolerance in human autoimmune thyroid disease. Mar 1998

4/6/6 09716948 98187354 PMID: 9526606
Preliminary studies with recombinant chorionic gonadotropin beta-subunit produced in Escherichia coli for use as an antigen in a birth control vaccine. Mar 1998

4/6/7 09511341 95156316 PMID: 7853237
Effects of thyroid hormone on fast- and slow-twitch skeletal muscles in young and old rats. Nov 15 1994

4/6/8 09419224 97479154 PMID: 9337806
Midlife periodic health exam in the primary care practice. Oct 1997

4/6/9 09175752 96429836 PMID: 8832986
Elevated serum prolactin or elevated prolactin/cortisol ratio are associated with autoimmune processes in systemic lupus erythematosus and other connective tissue diseases. Mar 1996

4/6/10 09073399 97025581 PMID: 8871760
The pathogenicity of spontaneously-occurring thyroglobulin-reactive T lymphocytes from BB/WOR rats. 1996

4/6/11 08491947 95229821 PMID: 7714099
Activation of T lymphocyte subsets by synthetic TSH receptor peptides and recombinant glutamate decarboxylase in autoimmune thyroid disease and insulin-dependent diabetes. Apr 1995

4/6/12 08365713 95213731 PMID: 7699384
Response of fast muscle innervation to hypothyroidism. Dec 1 1994

4/6/13 07938907 93390517 PMID: 8377760

[Evaluation on our procedure for autotransplantation of parathyroid glands by the intact-PTH] Aug 1993

4/6/14 07651909 93027039 PMID: 1408659
The time course of thyroid-hormone-induced changes in the isotonic and isometric properties of rat soleus muscle. Jul 1992

4/6/15 07355645 90368987 PMID: 2168443
Thyroid-stimulating antibody activity between different immunoglobulin G subclasses. Sep 1990

4/6/16 07344123 90192059 PMID: 2138281
The rate of tetanic relaxation is correlated with the density of calcium ATPase in the terminal cisternae of thyrotoxic skeletal muscle. Jan 1990

4/6/17 06343059 88056008 PMID: 3678672
Screening practices of family physicians: a comparison of STFM and AAFP members. Sep-Oct 1987

4/6/18 06175404 85196529 PMID: 3887952
Characteristics and bioefficacy of monoclonal antigenoadotropin releasing hormone antibody. Mar 1985

4/6/19 05788755 89067487 PMID: 2848890
Probing the normal and autoimmune B cell repertoire with Epstein-Barr virus. Frequency of B cells producing monoreactive high affinity autoantibodies in patients with Hashimoto's disease and systemic lupus erythematosus. Dec 15 1988

4/6/20 05639197 88043483 PMID: 3118511
Temporal analysis of diethylenetriamine neurotoxicity in rats and assessment of potential nonneural causes. Nov 1987

4/6/21 05514295 86248155 PMID: 2424791
Enhancement of antigenoadotropin response to the beta-subunit of ovine luteinizing hormone by carrier conjugation and combination with the beta-subunit of human chorionic gonadotropin. Jul 1986

4/6/22 05330203 90033436 PMID: 2806615
Antibody response and characteristics of antibodies in women immunized with three contraceptive vaccines inducing antibodies against human chorionic gonadotropin. Nov 1989

4/6/23 05287508 89332330 PMID: 2756340
Affinity purification of IgG subclasses and the distribution of thyroid auto-antibody reactivity in Hashimoto's thyroiditis. Jul 1989

4/6/24 05235053 88154038 PMID: 3257970
Microsomal antigen-reactive lymphocyte lines and clones derived from thyroid tissue of patients with Graves' disease. Apr 1988

4/6/25 04893804 80227723 PMID: 7391005
Interaction of fragments B and C of tetanus toxin with neural and thyroid membranes and with gangliosides. Jul 10 1980

4/6/26 04805868 82247718 PMID: 6179073
[Possible relation between the immune response and thymus-dependence of the immunizing antigens in children in an endemic goiter region] O vozmozhnoi svyazi mezhdu immunnym otvetom i timuszavisimosti u immunizirovannykh antigenov u detei v ochage zobnoi endemii. May-Jun 1982

4/6/27 04760092 84253338 PMID: 6331133
Gangliosides, the thyrotropin receptor, and autoimmune thyroid disease. 1984

4/6/28 04713242 80150428 PMID: 6244725
Thyrotropin receptors and gangliosides. 1980

4/6/29 04661467 84263155 PMID: 6378767
Defective regulation of the immune response to tetanus toxoid in Hashimoto's disease. Jul 1984

4/6/30 04655785 84158405 PMID: 6368201
Characterization of monoclonal antibody 3G5 and utilization of this antibody to immobilize pancreatic islet cell gangliosides in a solid phase radioassay. Apr 1984

4/6/31 03639753 82119926 PMID: 7328074
Thyroid function in tetanus. Jul 1981

4/6/32 03528268 79020956 PMID: 212049
Structure/function studies of receptors for thyrotropin and tetanus toxin: lipid modulation of effector binding to the glycoprotein receptor component. Jul 14 1978

4/6/33 03392187 79173087 PMID: 220221
Tetanus toxin and thyrotropin interactions with rat brain membrane preparations. May 25 1979

4/6/34 03389313 79127176 PMID: 217603

Tetanus toxin interactions with the thyroid: decreased toxin binding to membranes from a thyroid tumor with a thyrotropin receptor defect and in vivo stimulation of thyroid function. Mar 1978

4/6/35 03345587 76185875 PMID: 131619

[Permeability of sublingual mucosa to organic molecules. Limited role of sublingual absorption in aerosol vaccinations] La permeabilité de la

use sub-linguale aux molécules organiques. Les limites du rôle de la sub-linguale dans la vaccination par aérosols 1975

4/6/36 03008262 76102749 PMID: 813223

Isomunization against human chorionic gonadotropin with conjugates of processed beta-subunit of the hormone and tetanus toxoid. Jan 1976

4/6/37 02679440 77187913 PMID: 193853

4/7/2 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 10726207 20373526 PMID: 10916814

[Preventive activities in primary health care: identifying the agreement among evidence-based guidelines] Actividades preventivas en atención primaria: identificación de áreas de concordancia entre guías de práctica clínica basadas en la evidencia.

Gosalbes Soler V; Marquez Calderon S; Maiques Galan A; Latour Perez J; Bernal Delgado E; Puig Barbera J; Arranz Lazaro M

Unidad Docente de Medicina Familiar y Comunitaria, Valencia. gosalbesvic@gua.es
Medicina clínica (SPAIN) 2000, 114 Suppl 2 p88-92, ISSN 0025-7753 Journal Code: LTQ

Languages: SPANISH Document type: Journal Article Record type: Completed

BACKGROUND: The purpose of this article is to identify the agreement among evidence-based guidelines about recommendations on preventive activities in low risk adults. METHODS: For which we identified, from the 1996 US Task Force on Preventive Services Guide those preventives activities classified like A or B (recommendation in favour of provision) and like D or E (recommendation against provision), excepting those D and E recommendations based on descriptive studies or expert opinions. Both preventive activities aimed at pregnant women and children and those which are not applicable to our context were excluded. Selected preventive services were compared with the range of age, sex and periodicity in which agreement with the recommendations of American College of Physicians and Canadian Task Force on Preventive Services existed. RESULTS: We found the following agreements. Screening activities. In favour: screening for hypercholesterolemia, hypertension, breast cancer, colorectal cancer, uterine cervix cancer, rubella, visual and hearing impairment and problem drinking. Against: cancer of prostate, lung, bladder and thyroid, and asymptomatic bacteriuria. Counseling activities. In favour: smoking, motor-vehicles injuries, alcohol consumption, unintended pregnancy. Immunizations and quimioprophylaxis. In favour: Vaccines for influenza, tetanus-diphtheria, hepatitis B and measles-mumps-rubella. Postexposure prophylaxis to hepatitis A, hepatitis B, meningococcal, rabies and tetanus. CONCLUSIONS: We see then, that a high degree in agreement among the main guidelines exists; about the preventive activities to perform in Primary Health Services, nevertheless low fulfillment of certain preventive activities in Primary Health Services, different barriers for the accomplishment from these activities were described. Record Date Created: 20000919

4/7/5 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 09732112 98204670 PMID: 9545109

Induction of oral tolerance in human autoimmune thyroid disease.

Lee S; Scherberg N; DeGroot LJ

Department of Medicine, The University of Chicago, Illinois, USA.

Thyroid (UNITED STATES) Mar 1998, 8 (3) p229-34, ISSN 1050-7256 Journal Code: BJW
Languages: ENGLISH Document type: Clinical Trial; Journal Article; Randomized Controlled Trial Record type: Completed

Our laboratory has reported suppression of experimental autoimmune thyroiditis in mice by oral feeding with antigen. Based on these data, we considered it possible that oral feeding of animal thyroglobulin (TG) might induce tolerance to antigen in human autoimmune thyroid disease (AITD). Thirteen patients receiving thyroid hormone replacement with synthetic thyroxine (T4) (five patients with Graves' disease, treated with radioiodine 4 to 11 years ago and eight patients with Hashimoto's thyroiditis) were randomly assigned to a test group (switched to replacement with desiccated thyroid from porcine thyroids) and a control group (maintained on synthetic T4). Humoral and cellular immunologic parameters were evaluated in addition to clinical parameters before and every 3 months after the onset of study for a year. At the onset of study, there was no difference in clinical parameters, or humoral and cellular immunity to thyroid autoantigens, except a finding that one thyroid peroxidase (TPO) peptide (100 approximately 119) appeared to stimulate peripheral blood mononuclear cells (PBMC) during in vitro microproliferation assay more in the test group than control group ($p = 0.051$ by t test). Additionally, almost all of TPO and thyrotropin receptor extracellular domain (TSHR) peptides were slightly more stimulatory to PBMC from the test group than the control group, although this was not statistically significant. After treatment, all variables were analyzed at each time point between groups (t test), and also were analyzed over time in each group (analysis of variance, ANOVA). Among the clinical parameters, thyrotropin (TSH) levels were unchanged and equal. Total serum T4 levels ($p < 0.05$ at 6 and 12 months after treatment) and free thyroxine indices (FT4I) ($p < 0.05$ at all time points after treatment) were lower in the test group than the control group. This is an expected result of treatment with desiccated thyroid. We found no change over time nor any difference between groups at time points for titers of antibodies to thyroid autoantigens, ie, human TG, human TPO, and recombinant human TSHR from *Escherichia coli*. However, cellular immunity, measured by in vitro microproliferation of PBMC to peptides of TPO or TSHR, showed significant differences between groups. At 12 months, stimulatory indices (SI) of PBMC to six peptides, containing the indicated amino acids (764 approximately 95, 100 approximately 119, 110 approximately 129, 261 approximately 275, 441 approximately 448, 708 approximately 727) of 10 TPO peptides, and one peptide (145 approximately 163) of 14 TSHR peptides were lower in the test group than control group ($p < 0.05$). SI of PBMC to phytohemagglutinin, purified protein derivative from mycobacteria, and tetanus toxoid were not different between groups nor changed over time in any group. In conclusion, treating patients with AITD with an antigen related to the autoantigen TG did not produce changes in humoral immunity parameters, while cellular immunity to certain peptides were apparently suppressed. While the results are both surprising and intriguing, we need more evidence to justify the use of autoantigen as a form of immunospecific therapy in patients with AITD. Record Date Created: 19980515

Tetanus toxin interactions with thyroid plasma membranes. Implications for structure and function of tetanus toxin receptors and potential pathophysiological significance. Jun 25 1977

4/6/38 01235275 70276257 PMID: 4916539

Immunosuppressive therapy for the eye changes of Graves' disease. Sep 1970

4/7/7 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 09511341 95156316 PMID: 7853237

Effects of thyroid hormone on fast- and slow-twitch skeletal muscles in young and old rats.

Larsson L; Li X; Teresi A; Salvati G

Department of Clinical Neurophysiology, Karolinska Hospital, Stockholm, Sweden.

Journal of physiology (ENGLAND) Nov 15 1994, 481 (Pt 1) p149-61, ISSN 0022-3751

Journal Code: JQV Languages: ENGLISH Document type: Journal Article Record type: Completed

1. The effects of 4 weeks of thyroid hormone treatment on contractile, enzyme-histochemical and morphometric properties and on the myosin isoform composition were compared in the slow-twitch soleus and the fast-twitch extensor digitorum longus (EDL) muscle in young (3-6 months) and old (20-24 months) male rats. 2. In the soleus of untreated controls, contraction and half-relaxation times of the isometric twitch increased by 19-32% with age. The change in contractile properties was paralleled by an age-related increase in the proportions of type I fibres and type I myosin heavy chain (MHC) and slow myosin light chain (MLC) isoforms. 3. In the EDL of controls, contraction and half-relaxation times were significantly prolonged (21-38%) in the post-tetanus twitch in the old animals. No significant age-related changes were observed in enzyme-histochemical fibre-type proportions, although the number of fibres expressing both type IIA and IIB MHCs and of fibres expressing slow MLC isoforms was increased in the old animals. 4. Serum 3,5,3',5'-tetraiodothyronine (T4) levels were lower (34%) in the old animals, but the primary byproduct of T4, 3,5,3'-triiodothyronine (T3), did not differ between young and old animals. 5. The effects of 4 weeks of thyroid hormone treatment were highly muscle specific, and were more pronounced in soleus than in EDL, irrespective of animal age. In the soleus, this treatment shortened the contraction and half-relaxation times by 35-57% and decreased the number of type I fibres by 66-77% in both young and old animals. In EDL, thyroid hormone treatment significantly shortened the contraction time by 24%, but the change was restricted to the old animals. 6. In conclusion, the ability of skeletal muscle to respond to thyroid hormone treatment was not impaired in old age and the age-related changes in speed of contraction and enzyme-histochemical properties and myosin isoform compositions were diminished after thyroid hormone treatment in both the soleus and EDL. Record Date Created: 19950314

4/7/12 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 08365713 95213731 PMID: 7699384

Response of fast muscle innervation to hypothyroidism.

Cuppini R; Sartini S; Ambrogini P; Gallo G

Istituto di Anatomia e Fisiologia, Università di Urbino, Italy.

Journal of the neurological sciences (NETHERLANDS) Dec 1 1994, 127 (1) p107-13, ISSN 0022-510X Journal Code: JBJ Languages: ENGLISH Document type: Journal Article Record type: Completed

The early period of motor innervation development is characterized by multiple innervation of muscle cells. This transitory state in rat extensor digitorum longus (edl) muscle is normally concluded at weaning when a 1:1 ratio between nerve endings and muscle cells is reached. Motor innervation of edl muscle in rats made hypothyroid after weaning was studied in three ways: electrophysiology (intracellular recordings of muscle postsynaptic potentials) was carried out to study neuromuscular transmission; silver impregnation of terminal axons to observe sprouting; force production in twitch and tetanus following direct muscle stimulation and nerve stimulation. A number of multiply innervated muscle cells was found in hypothyroid rats following two months of treatment. This finding seems to be related to the appearance of nodal sprouting in motor axons. No sign of denervated end-plates was found. Twitch and tetanus tension were smaller than in controls, but they were bigger when referred to unitary muscle mass. Time course of twitch, particularly half relaxation, was slowed in muscles of hypothyroid rats. These findings suggest that plastic processes occur in muscle innervation of rats made hypothyroid after weaning. Therefore, thyroid hormones play a role in stabilizing motor innervation not only during development, but also in adults. Record Date Created: 19950428

4/7/13 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 07938907 93390517 PMID: 8377760

[Evaluation on our procedure for autotransplantation of parathyroid glands by the intact-PTH]

Ueda M; Funahashi H; Sato Y; Kato M; Takagi H

Second Department of Surgery, Nagoya University, School of Medicine, Japan.

Nippon Geka Gakkai zasshi (JAPAN) Aug 1993, 94 (8) p840-6, ISSN 0301-4894 Journal Code: NGG Languages: JAPANESE Document type: Journal Article Record type: Completed
In thyroid cancer surgery since 1978 we have made parathyroidal autotransplantation which has been to resect the entire parathyroid, cut fine to mud and autotransplant into greater pectoral muscle. To evaluate our procedure and consider the mechanism of hypocalcemia and tetanus following surgery, we examined the intact-PTH recovered to about 80% of its preoperative value on 14-postoperative-days and thereafter remained almost constant. It remained below the sensitivity of measurement from immediately to 3-postoperative-days, and with our procedure, parathyroid was considered to be totally resected. Comparison between 2 (6 cases) and more autotransplanted glands (11 cases) revealed that the former showed slightly later functional recovery, but recovered to the almost same extent as the other 21-postoperative-days. Comparison between the group requiring calcium supplement therapy (12 cases) and the otherwise (5 cases) revealed almost the same course of recovery. Thus, our procedure seems to enable us to make satisfactory functional preservation and expect functional recovery by 2-gland

autotransplantation at least. The supplement therapy was considered to be detrimental to the take of autotransplanted glands and functional recovery, and no correlation was noted between the onset of tetanus and intact-PTH. Record Date Created: 19931020

4/7/17 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 06343059 88056008 PMID: 3678672

Screening practices of family physicians: a comparison of STFM and AAFP members.

Resnicow K; Schorow M; Bloom HG; Massad R; Coll-Barth M

American Health Foundation, New York City.

Family medicine (UNITED STATES) Sep-Oct 1987, 19 (5) p341-5, ISSN 0742-3225

Journal Code: FAL Languages: ENGLISH Document type: Journal Article Record type: Completed

The screening practices of 146 members of the Society of Teachers of Family Medicine (STFM) and 129 members of the American Academy of Family Physicians (AAFP) are compared. The screening practices of physicians from the two organizations were generally similar for psychosocial and behavioral problems, many forms of cancer, and numerous other conditions considered for inclusion in the routine periodic screening of asymptomatic individuals. However, for numerous diseases and tests, the screening practices of physicians from the two groups were significantly different. AAFP physicians were more likely to screen for lung and skin cancer, thyroid dysfunction, diabetes, and anemia, AAFP physicians were more likely to utilize chest x-ray, ECG, urinalysis, and SMA 6/12. STFM physicians were more likely to perform gonococcal culture and tetanus-diphtheria immunization as well as to inquire about seat belt use. Three variables were found to predict physician screening practices as well as to account for the differences found between physicians drawn from the two organizations: completion of a residency in family medicine, year of graduation from medical school, and number of patients seen per week. Physicians reported practices were compared with recommendations in the major critical reviews: Frame and Carlson, Breslow and Sommers, the Canadian Task Force, and the American Cancer Society. For a number of tests and diseases physicians' reported practices were divergent with recent recommendations. Record Date Created: 19871221

4/7/20 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 05639197 88043483 PMID: 3118511

Temporal analysis of dithiobiuret neurotoxicity in rats and assessment of potential nonneural causes.

Williams KD; Miller MS; Boysen BG; Peterson RE

School of Pharmacy, University of Wisconsin, Madison 53706.

Toxicology and applied pharmacology (UNITED STATES) Nov 1987, 91 (2) p212-21, ISSN 0041-008X Journal Code: VWO Contract/Grant No.: 1-K04-ES00098, ES, NIEHS; ES01906, ES, NIEHS Languages: ENGLISH Document type: Journal Article Record type: Completed

To evaluate the hypothesis that depressed neuromuscular transmission causes dithiobiuret (DTB)-induced muscle weakness in rats, the temporal development of impaired treadmill performance and deficits in the nerve-elicited muscle contractions were compared during daily treatment with the toxicant (DTB, 1 mg/kg/day X 6 days). Diminished treadmill test performance after 4 days of treatment marked the initial detection of impaired motor function. At this time fading (loss of tension during tetanus) of gastrocnemius contractions elicited in response to 100-Hz sciatic nerve stimulation occurred in DTB-treated rats but not in controls. After 5 and 6 days of treatment, treadmill failure became complete, tetanic fade worsened dramatically, and peak contractile tension measured during trains of nerve stimulation (10-250 Hz) decreased progressively. Appearing by Day 6 were marked body weight loss, dehydration, hypothermia, and a depression in serum concentrations of thyroid hormones. Total oxygen content of the blood was not reduced at any time during treatment, and serum concentrations of glucose, sodium, potassium, calcium, chloride, and phosphorus in DTB-treated rats on Day 6 were similar to those of control animals. Therefore, hypoxia, hypoglycemia, or a serum electrolyte imbalance do not initiate or modulate the neuromuscular toxicity. Light microscopic evaluation of liver, kidney, lung, thyroid, and other organs in intoxicated rats was unremarkable and in skeletal muscles and selected sites of brain, spinal cord, and sciatic nerve no morphologically significant lesions were observed. Even when DTB-intoxicated rats were maintained in a state of flaccid muscle weakness for 5 continuous days, peripheral nerve lesions proximal to the intramuscular nerves were not detected. Thus, depressed neuromuscular transmission appears to be the primary cause of the flaccid muscle weakness and no evidence was obtained that nonneural effects of DTB initiate or modulate this effect. Record Date Created: 19871211

4/7/24 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 05235053 88154038 PMID: 3257970

Microsomal antigen-reactive lymphocyte lines and clones derived from thyroid tissue of patients with Graves' disease.

Fisfalen ME; DeGroot LJ; Quintans J; Franklin WA; Soltani K

Department of Medicine, University of Chicago, Illinois 60637.

Journal of clinical endocrinology and metabolism (UNITED STATES) Apr 1988, 66 (4)

p776-84, ISSN 0021-972X Journal Code: HRB Contract/Grant No.: AM-13377, AM, NIADDK; CA19589, CA, NCI; DK-27384, DK, NIDDK

Languages: ENGLISH Document type: Journal Article Record type: Completed

Thyroid mononuclear cells (TMC) were maintained in long term cocultures with thyroid fibroblasts and thyroid epithelial cells from patients with Graves' disease, using medium supplemented with thyroid microsomal antigen (McAg) and IL-2. The TMC consisted predominantly of T4+ (CD4+, helper) and, to a lesser extent, T8+ (CD8+, cytotoxic/suppressor) lymphocytes, with a small number of macrophages and natural killer cells. The average T4+ to T8+ ratio was 3.2. From these cultures we obtained thyroid T cell lines and clones reactive to thyroid antigens. T cell lines were tested in a microproliferation assay using thyroglobulin (Tg), McAg, tetanus toxoid, and IL-2. Of 14 lines from 6 patients, 2 proliferated in response to McAg when TMC plus thyroid fibroblasts were used as antigen-presenting cells. Clones of thyroid lymphocytes were obtained by culturing cells at limiting dilution with IL-2, McAg, and different types of autologous accessory cells.

Peripheral blood mononuclear cells and skin fibroblasts provided the best source of accessory cells, allowing near 100% cloning efficiency. Of 26 clones tested, 6 recognized McAg, 2 were Tg reactive, and 3 were autoreactive. All phenotyped clones were of the T4+ phenotype. Our method results in production of thyroid T cell lines and clones. The fibroblasts probably provided growth factors and/or collaborated with peripheral blood mononuclear cells as antigen-presenting cells. These lines and clones from patients with Graves' disease were predominantly helper T cells, in contrast to the previously demonstrated cytotoxic/suppressor cell predominance in cells from patients with Hashimoto's thyroiditis. This difference in cell function may help explain the differing clinical courses of these two closely related autoimmune thyroid diseases. The availability of long term microsomal antigen-specific T cell clones should allow careful analysis of the role these cells play in thyroid autoimmunity. Record Date Created: 19880419

4/7/25 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 04893804 80227723 PMID: 7391005

Interaction of fragments B and C of tetanus toxin with neural and thyroid membranes and with gangliosides.

Morris NP; Consiglio E; Kohn LD; Habis WH; Hardegree MC; Helting TB

Journal of biological chemistry (UNITED STATES) Jul 10 1980, 255 (13) p6071-6, ISSN 0021-9258 Journal Code: HIV

Languages: ENGLISH Document type: Journal Article Record type: Completed Record Date Created: 19800928

4/7/27 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 04760092 84253338 PMID: 6331133

Gangliosides, the thyrotropin receptor, and autoimmune thyroid disease.

Lacetti P; Tombaccini D; Aloj S; Grollman EF; Kohn LD

Advances in experimental medicine and biology (UNITED STATES) 1984, 174 p355-67, ISSN 0065-2598 Journal Code: 2LU Languages: ENGLISH Document type: Journal Article Record type: Completed

The thyrotropin (TSH) receptor has been proposed to be composed of a membrane glycoprotein and a membrane ganglioside, the former important in high affinity recognition, the latter vital for message coupling to the adenylate cyclase system. The present study used two approaches, formation of antireceptor monoclonal antibodies and reconstitution, to validate the model and further examine the role of the ganglioside. Three kinds of monoclonal antireceptor antibodies are defined. One group which inhibits TSH binding and TSH functions, i.e., TSH-stimulated adenylate cyclase activity, iodide uptake, and thyroid hormone release, is shown to be directed against the glycoprotein component of the receptor. The second group includes antibodies which mimic TSH in all stimulatory actions, are competitive agonists of TSH, are equivalent to thyroid stimulating antibodies in the sera of patients with Graves' disease, and are directed against the ganglioside component of the receptor. These stimulating monoclonal antibodies are directed against a minor ganglioside membrane component which fractionates as a disialoganglioside. When this ganglioside is incorporated into 1-8 thyroid cells which have a correlated ganglioside deficiency and TSH receptor defect, reconstitution of TSH stimulated adenylate cyclase activity occurs. Whereas the first group of antibodies inhibits TSH-stimulated function, they do not inhibit the stimulatory antibodies which mimic TSH, an observation consistent with the 2 component hypothesis of the receptor model. The third group of antibodies have a mix of properties from the first two groups and suggests that the TSH receptor in situ is an actual complex of the two components or that there are common carbohydrate determinants in the functional sites of each receptor component. Implications of a TSH receptor structure in which its ganglioside and glycoprotein components are in equilibrium with pools of free components and, in turn, components important for cholera toxin, tetanus toxin and interferon receptors are discussed. In regard to the pathogenesis of Graves' disease, the data indicate that thyroid stimulating autoantibodies are autoimmune equivalents of cholera toxin with respect to the importance of ganglioside function. Since antidiotype studies of antibodies against TSH confirm a structural relationship between receptors for thyrotropin, cholera toxin, and thyroid stimulating autoantibodies, the data establish an unequivocal role for the ganglioside in TSH receptor structure which facilitates interpretation of in vitro experiments aimed at understanding the mechanism of ganglioside-ligand interactions. Record Date Created: 19840823

4/7/28 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 04713242 80150428 PMID: 6244725

Thyrotropin receptors and gangliosides.

Kohn LD; Consiglio E; DeWolf MJ; Grollman EF; Ledley FD; Lee G; Morris NP

Advances in experimental medicine and biology (UNITED STATES) 1980, 125 p487-503, ISSN 0065-2598 Journal Code: 2LU Languages: ENGLISH Document type: Journal Article; Review Record type: Completed (44 Refs.) Record Date Created: 19800530

4/7/29 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 04661467 84263155 PMID: 6378767

Defective regulation of the immune response to tetanus toxoid in Hashimoto's disease.

Fawcett J; Hutton C; McLachlan SM; Clark F; Rees Smith B

Immunology (ENGLAND) Jul 1984, 52 (3) p525-8, ISSN 0019-2805 Journal Code: GH7

Languages: ENGLISH Document type: Journal Article Record type: Completed

The humoral immune response to tetanus toxoid has been studied in patients with Hashimoto's disease. Although the magnitude of the response was similar to that observed in normal subjects, the Hashimoto patients demonstrated an inability to regulate their levels of tetanus toxoid antibody. This apparent defect in the control of antibody synthesis may be an important factor in both the initiation and perpetuation of autoimmune thyroid disease. Record Date Created: 19840829

4/7/31 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 03639753 82119926 PMID: 7328074

Thyroid function in tetanus.
Dastur FD; Shah J; Awatramani V; Shah JM; Pardiwalla BS; Nair KG; Mehta MN; Desai KB
Journal of the Association of Physicians of India (INDIA) Jul 1981, 29 (7) p519-25, ISSN
0004-5772 Journal Code: HG7 Languages: ENGLISH Document type: Journal Article
Record type: Completed Record Date Created: 19820412

4/5/34 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
03389313 79127176 PMID: 217603

Tetanus toxin interactions with the thyroid: decreased toxin binding to membranes from a thyroid tumor with a thyrotropin receptor defect and in vivo stimulation of thyroid function.
Habig WH; Grollman EF; Ledley FD; Meldolesi MF; Aloj SM; Hardegree MC; Kohn LD
Endocrinology (UNITED STATES) Mar 1978, 102 (3) p844-51, ISSN 0013-7227 Journal
Code: EGZ Languages: ENGLISH Document type: Journal Article Record type: Completed
Normal rat thyroid membranes adsorb neurotoxicity when incubated with purified tetanus toxin.
Membranes from a rat thyroid tumor with a thyrotropin receptor defect adsorb very little
neurotoxicity when similarly evaluated. This inability of the tumor membranes to adsorb
neurotoxicity is correlated with a defect in their ability to bind both 125I-labeled tetanus toxin and
[125I]iodothyrotropin. The effect of tetanus toxin on the release of radioiodine from the thyroids of
appropriately prepared mice has been measured by adapting methods used for the bioassay of
thyrotropin. One minimum lethal dose of tetanus toxin given sc caused a significant release of
radioiodine into the blood of mice 48 h after injection. In mice subjected to the stress of prior
bleedings or anesthesia, the release of radioiodine from the thyroid by tetanus toxin was
accelerated, i.e., the increase in blood radioiodine could be measured 24 h after injection. These
results again suggest that tetanus toxin may interact with thyrotropin receptors on thyroid plasma
membranes. The "sympathetic overactivity syndrome" seen in some patients with tetanus and the
7/6/1 09732112 98204670 PMID: 9545109
Antigenic determinants on human choriogonadotropin alpha-subunit. II.
Induction of oral tolerance in human autoimmune thyroid disease. Mar 1998
Immunochemical mapping by a monoclonal antipeptide antibody. Jul 25 1988

7/6/2 09408392 98018742 PMID: 9376076
TCR vbeta usage of TSH receptor-specific CD4+ T cells in Graves' disease patients and healthy humans. Oct 1997

7/6/3 08491947 95229821 PMID: 7714099
Activation of T lymphocyte subsets by synthetic TSH receptor peptides and recombinant glutamate decarboxylase in autoimmune thyroid disease and insulin-dependent diabetes. Apr 1995

7/6/4 07355645 90368987 PMID: 2168443
Thyroid-stimulating antibody activity between different immunoglobulin G subclasses. Sep 1990

7/6/5 07320865 91048427 PMID: 2122534
Thyrotropin-releasing hormone has profound presynaptic action on cultured spinal cord neurons. 1990

7/6/6 06175404 85196529 PMID: 3887952
Characteristics and bioefficacy of monoclonal antigonadotropin releasing hormone antibody. Mar 1985

7/6/7 05550007 88273140 PMID: 2455715

7/7/12 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
03639753 82119926 PMID: 7328074
Thyroid function in tetanus.

Dastur FD; Shah J; Awatramani V; Shah JM; Pardiwalla BS; Nair KG; Mehta MN; Desai KB
Journal of the Association of Physicians of India (INDIA) Jul 1981, 29 (7) p519-25, ISSN
0004-5772 Journal Code: HG7 Languages: ENGLISH Document type: Journal Article
Record type: Completed Record Date Created: 19820412

7/7/15 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
03389313 79127176 PMID: 217603

Tetanus toxin interactions with the thyroid: decreased toxin binding to membranes from a thyroid tumor with a thyrotropin receptor defect and in vivo stimulation of thyroid function.
Habig WH; Grollman EF; Ledley FD; Meldolesi MF; Aloj SM; Hardegree MC; Kohn LD
Endocrinology (UNITED STATES) Mar 1978, 102 (3) p844-51, ISSN 0013-7227 Journal
Code: EGZ Languages: ENGLISH Document type: Journal Article Record type: Completed
Normal rat thyroid membranes adsorb neurotoxicity when incubated with purified tetanus toxin.
Membranes from a rat thyroid tumor with a thyrotropin receptor defect adsorb very little

10/6/1 04722792 81133651 PMID: 6258596
Demonstration and characterization of partial glyceride specific lipases in pig thyroid plasma membranes. Nov 28 1980

10/6/2 04661467 84263155 PMID: 6378767
Defective regulation of the immune response to tetanus toxoid in Hashimoto's disease. Jul 1984

10/6/3 02675218 77138183 PMID: 191247

10/5/4 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
01695705 67139798 PMID: 4960830
Action of phospholipase C on the thyroid. Abolition of the response to thyroid-stimulating hormone.
Macchia V; Pastan I

Effects of concanavalin A and neuraminidase on cyclic AMP levels and 14C-1-glucose oxidation in dog thyroid slices. Aug 1976

10/6/4 01695705 67139798 PMID: 4960830
Action of phospholipase C on the thyroid. Abolition of the response to thyroid-stimulating hormone. Apr 25 1967

10/6/5 01512434-70106909 PMID: 4312983
Role of lecithin in the mechanism of TSH action. Apr 1970

syndrome characterized as "thyroid storm" in patients with Graves' disease are discussed as they may relate to these observations. Record Date Created: 19790516

Tags: Animal
Descriptors: Protirelin-metabolism-ME; *Receptors, Cell Surface-metabolism-ME; *Tetanus Toxin-metabolism-ME; *Thyroid Gland-metabolism-ME; *Thyroid Neoplasms-metabolism-ME; Cell Membrane-metabolism-ME; Kinetics; Rats
CAS Registry No.: 0 (Receptors, Cell Surface); 0 (Tetanus Toxin); 24305-27-9 (Protirelin)

5/6/1 02255653 69254786 PMID: 4895160
[Various findings on the processes of natural detoxication of the body during infectious processes]
Nekotorye dannye o protsessakh estestvennoi detoksikatsii organizma pri infektsionnykh protsessakh. Oct 1967

5/5/2 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
00882933 71059144 PMID: 5487765
Clostridium septicum infection of the thyroid gland.
Warren CP; Mason BJ
Postgraduate medical journal (ENGLAND) Sep 1970, 46 (539) p586-8, ISSN 0032-5473
Journal Code: PFX
Languages: ENGLISH Document type: Journal Article Record type: Completed Record Date Created: 19710204
Tags: Human; Male
Descriptors: Clostridium Infections-complications-CO; *Thyroiditis-etiology-ET; Adenocarcinoma-complications-CO; Clostridium-isolation and purification-IP; Colonic Neoplasms-complications-CO; Middle Age; Thyroiditis-microbiology-MI

7/6/14 03392187 79173087 PMID: 220221
Tetanus toxin and thyrotropin interactions with rat brain membrane preparations. May 25 1979

7/6/15 03389313 79127176 PMID: 217603
Tetanus toxin interactions with the thyroid: decreased toxin binding to membranes from a thyroid tumor with a thyrotropin receptor defect and in vivo stimulation of thyroid function. Mar 1978

7/6/16 03289145 79015004 PMID: 358455
Membrane receptors for interferon. 1977

7/6/17 02869495 76091246 PMID: 1245125
Immunological reactivity of antibodies produced by Pr-beta-HCG-TT with different hormones. Feb 1976

7/6/18 02679440 77187913 PMID: 193853
Tetanus toxin interactions with thyroid plasma membranes. Implications for structure and function of tetanus toxin receptors and potential pathophysiological significance. Jun 25 1977

7/6/19 02618964 78053767 PMID: 337386
[Biosynthesis and function of gangliosides (author's transl)] 1977

neurotoxicity when similarly evaluated. This inability of the tumor membranes to adsorb neurotoxicity is correlated with a defect in their ability to bind both 125I-labeled tetanus toxin and [125I]iodothyrotropin. The effect of tetanus toxin on the release of radioiodine from the thyroids of appropriately prepared mice has been measured by adapting methods used for the bioassay of thyrotropin. One minimum lethal dose of tetanus toxin given sc caused a significant release of radioiodine into the blood of mice 48 h after injection. In mice subjected to the stress of prior bleedings or anesthesia, the release of radioiodine from the thyroid by tetanus toxin was accelerated, i.e., the increase in blood radioiodine could be measured 24 h after injection. These results again suggest that tetanus toxin may interact with thyrotropin receptors on thyroid plasma membranes. The "sympathetic overactivity syndrome" seen in some patients with tetanus and the syndrome characterized as "thyroid storm" in patients with Graves' disease are discussed as they may relate to these observations. Record Date Created: 19790516

Tags: Animal
Descriptors: Protirelin-metabolism-ME; *Receptors, Cell Surface-metabolism-ME; *Tetanus Toxin-metabolism-ME; *Thyroid Gland-metabolism-ME; *Thyroid Neoplasms-metabolism-ME; Cell Membrane-metabolism-ME; Kinetics; Rats
CAS Registry No.: 0 (Receptors, Cell Surface); 0 (Tetanus Toxin); 24305-27-9 (Protirelin)

10/6/6 01502036 68318489 PMID: 4298230
Effect of sphingomyelinase from Clostridium perfringens on the metabolic activity and phospholipid composition of thyroid slices. Jul 1 1968

10/6/7 01496873 67221632 PMID: 4292226
The purification and properties of a thyroid-stimulating factor isolated from Clostridium perfringens. Aug 25 1967

Journal of biological chemistry (UNITED STATES) Apr 25 1967, 242 (8) p1864-9, ISSN 0021-9258 Journal Code: HIV
Languages: ENGLISH Document type: Journal Article Record type: Completed Subfile: INDEX MEDICUS
Tags: Animal; In Vitro

Descriptors: Phospholipases--pharmacology--PD; *Thyroid Gland--drug--DE; *Thyrotropin--pharmacology--PD; Acetylcholine--pharmacology--PD; Clostridium--enzymes--EN; Dogs; Edetic Acid--pharmacology--PD; Glucose--metabolism--ME; Neuraminidase--metabolism--ME
 CAS Registry No.: 50-99-7 (Glucose); 51-84-3 (Acetylcholine); 60-00-4 (Edetic Acid); 9002-71-5 (Thyrotropin) Enzyme No.: EC 3.1.- (Phospholipases); EC 3.2.1.18 (Neuraminidase) Record Date Created: 19670706

10/7/77 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
 01496873 67221632 PMID: 4292226

11/6/1 11237950 21017795 PMID: 11145348
 Spectroscopic investigation of calcium binding sites in the neurotoxin vipoxin and its components--relation with the X-ray structure. Dec 2000

11/6/2 10955619 21079817 PMID: 11212293
 Modulation of phospholipase A2 activity generated by molecular evolution. Oct 30 1999

11/6/3 10938101 20556627 PMID: 11102474
 alpha-Latrotoxin releases calcium in frog motor nerve terminals. Dec 1 2000

11/6/4 10921630 20458278 PMID: 11005268
 Sexual dimorphism in the hypothalamo-pituitary-adrenal (HPA) axis and TNFalpha responses to phospholipase A2-related neurotoxin (from *Crotalus durissus terrificus*) challenge. Jul-Aug 2000

11/6/5 10902774 20534643 PMID: 11080683
 Specific physiological roles of cytosolic phospholipase A(2) as defined by gene knockouts. Oct 31 2000

11/6/6 10742898 98055712 PMID: 9395334
 Characterization and evolution of a gene encoding a Trimeresurus flavoviridis serum protein that inhibits basic phospholipase A2 isozymes in the snake's venom. Nov 1 1997

11/6/7 10732492 20363696 PMID: 10903499
 Genetic organization of A chain and B chain of beta-bungarotoxin from Taiwan banded krait (*Bungarus multicinctus*). A chain genes and B chain genes do not share a common origin. Aug 2000

11/6/8 10610705 20223270 PMID: 10759860
 Combining phage display and molecular modeling to map the epitope of a neutralizing antitoxin antibody. Apr 2000

11/6/9 10561005 20200644 PMID: 10736105
 Neutralizing human anti crotoxin scFv isolated from a nonimmunized phage library. Apr 2000

11/6/10 10518268 20062793 PMID: 10594896
 Phospholipase A2 and arachidonic acid-mediated mechanism of neuroexocytosis: a possible target of botulinum neurotoxin A other than SNAP-25. Dec 1999

11/6/11 10516379 20133101 PMID: 10666574
 Structure of agkistrotoxin in an orthorhombic crystal form with six molecules per asymmetric unit. Dec 1999

11/6/12 10515269 20143523 PMID: 10677596
 Possible involvement of cytosolic phospholipase A(2) in cell death induced by 1-methyl-4-phenylpyridinium ion, a dopaminergic neurotoxin, in GH3 cells. Feb 14 2000

11/6/13 10379076 99447141 PMID: 10519652
 Studies on the specificity of CNF, a phospholipase A2 inhibitor isolated from the blood plasma of the South American rattlesnake (*Crotalus durissus terrificus*). I. Interaction with PLA2 from *Lachesis muta muta* snake venom. Dec 1999

11/6/14 10337859 99311688 PMID: 10384962
 Molecular mimicry between a monoclonal antibody and one subunit of crotoxin, a heterodimeric phospholipase A2 neurotoxin. May 1999

11/6/15 10312466 98406156 PMID: 9733637
 Crystal structure of agkistrotoxin, a phospholipase A2-type presynaptic neurotoxin from *Agkistrodon halys pallas*. Sep 11 1998

11/6/16 10164153 99234372 PMID: 10216316
 Isolation, purification, crystallization and preliminary X-ray analysis of beta 1-bungarotoxin from *Bungarus caeruleus* (Indian common krait). May 1999

11/6/17 10155746 99177348 PMID: 10077532
 Postsynaptic alpha-neurotoxin gene of the spitting cobra, *Naja naja sputatrix*: structure, organization, and phylogenetic analysis. Mar-1999

11/6/18 10151808 99233142 PMID: 10218774
 Acute application of the tricyclic antidepressant desipramine presynaptically stimulates the exocytosis of glutamate in the hippocampus. Mar 1999

11/6/19 10072498 99131241 PMID: 9932449

The purification and properties of d-stimulating factor isolated from *Clostridium perfringens*.

Macchia V; Bates RW; Pastan I
 Journal of biological chemistry (UNITED STATES) Aug 25 1967, 242 (16) p3726-30, ISSN 0021-9258 Journal Code: HIV
 Languages: ENGLISH Document type: Journal Article Record type: Completed Record Date Created: 19671022

The neuromessenger platelet-activating factor in plasticity and neurodegeneration. 1998

11/6/20 10027968 99117402 PMID: 9918538
 Characterization of pardaxin-induced dopamine release from pheochromocytoma cells: role of calcium and eicosanoids. Feb 1999

11/6/21 09997106 99081796 PMID: 9864269
 Pardaxin, a new pharmacological tool to stimulate the arachidonic acid cascade in PC12 cells. Dec 1998

11/6/22 09907469 98368852 PMID: 9705154
 A monoclonal antibody directed against the non-toxic subunit of a dimeric phospholipase A2 neurotoxin, crotoxin, neutralizes its toxicity. Jul 1998

11/6/23 09899508 98364102 PMID: 9698946
 Spectroscopic properties and stability of the neurotoxic complex. Vipoxin and its components. Aug 1998

11/6/24 09818845 98267642 PMID: 9604283
 Some pharmacological studies of venom from the inland taipan (*Oxyuranus microlepidotus*). Jan 1998

11/6/25 09786197 98303724 PMID: 9637731
 Purification, characterization, and amino acid sequence determination of acanthins, potent inhibitors of platelet aggregation from *Acanthopis antarcticus* (common death adder) venom. Jun 15 1998

11/6/26 09664799 98106963 PMID: 9444558
 Effect of 1-methyl-4-phenylpyridinium (MPP+) on mitochondrial membrane potential in cerebellar neurons: interaction with the NMDA receptor. 1997

11/6/27 09647820 98065610 PMID: 9402043
 Presynaptic snake beta-neurotoxins produce tetanic fade and endplate potential run-down during neuromuscular blockade in mouse diaphragm. Nov 1997

11/6/28 09638997 98061472 PMID: 9399104
 Mechanisms involved in activation of human eosinophil exocytosis by substance P: an in vitro model of sensory neuroimmunomodulation. Aug-Dec 1997

11/6/29 09629054 98086776 PMID: 9417893
 Inferring species trees from gene trees: a phylogenetic analysis of the Elapidae (Serpentes) based on the amino acid sequences of venom proteins. Dec 1997

11/6/30 09587245 97365572 PMID: 9222399
 Single-step purification of crotoxin and crotoxin from *Crotalus durissus terrificus* venom using preparative isoelectric focusing. Jan 1997

11/6/31 09532541 97450547 PMID: 9305535
 The histamine releasers croptamine, protamine and compound 48/80 activate specific proteases and phospholipases A2. Sep 1997

11/6/32 09317983 97179777 PMID: 9028013
 Identification of phospholipase A2 and neurotoxic activities in the venom of the New Guinean small-eyed snake (*Micropechis ikaheka*). Jan 1997

11/6/33 09277700 97215815 PMID: 9062105
 Neurotoxic phospholipases A2 ammodytoxin and crotoxin bind to distinct high-affinity protein acceptors in *Torpedo marmorata* electric organ. Mar 11 1997

11/6/34 09196094 97091771 PMID: 8937437
 Differential effects of snake venom phospholipase A2 neurotoxin (beta-bungarotoxin) and enzyme (*Naja naja atra*) on protein kinases. Oct 25 1996

11/6/35 09174482 97179741 PMID: 9027977
 Mechanism of action of beta-bungarotoxin, a presynaptically acting phospholipase A2 neurotoxin: its effect on protein phosphorylation in rat brain synaptosomes. Nov-Dec 1996

11/6/36 09045162 97085325 PMID: 8931466
 Two components of glutamate exocytosis differentially affected by presynaptic modulation. Dec 1996

11/6/37 08961877 96353154 PMID: 8748373

Rapid activity-dependent sprouting of optic fibers into a local area denervated by application of beta-bungarotoxin in goldfish tectum. Jan 1996

11/6/38 08923268 96248443 PMID: 8665942
 Characterization of alpha-neurotoxin and phospholipase A2 activities from *Micrurus* venoms. Determination of the amino acid sequence and receptor-binding ability of the major alpha-neurotoxin from *Micrurus nigrocinctus nigrocinctus*. May 15 1996

11/6/39 08777080 95406914 PMID: 7676466
 beta-Bungarotoxin blocks phorbol ester-stimulated phosphorylation of MARCKS, GAP-43 and synapsin I in rat brain synaptosomes. Jun 1995

11/6/40 08718843 96172663 PMID: 8588211
 Amino acid sequences of a heterodimeric neurotoxin from the venom of the false horned viper (*Pseudocerastes fieldi*). Jul 1995

11/6/41 08716806 96131881 PMID: 8590005
 Structure of beta 2-bungarotoxin: potassium channel binding by Kunitz modules and targeted phospholipase action. Oct 15 1995

11/6/42 08662024 96067607 PMID: 7487947
 Molecular cloning and characterization of a neurotoxic phospholipase A2 from the venom of Taiwan habu (*Trimeresurus mucrosquamatus*). Nov 1 1995

11/6/43 08632072 96018127 PMID: 7570629
 Ammodytoxin A acceptor in bovine brain synaptic membranes. Apr 1995

11/6/44 08195967 94307256 PMID: 8033889
 The origin of the diversity of crotoxin isoforms in the venom of *Crotalus durissus terrificus*. Jul 1 1994

11/6/45 08145101 94253151 PMID: 8195214
 A phospholipase A2 inhibitor from the plasma of the South American rattlesnake (*Crotalus durissus terrificus*). Protein structure, genomic structure, and mechanism of action. Jun 3 1994

11/6/46 08140009 94223544 PMID: 8169833
 Inhibition of vacuolar adenosine triphosphatase antagonizes the effects of clostridial neurotoxins but not phospholipase A2 neurotoxins. Apr 1994

11/6/47 08126117 94182951 PMID: 8135548
 Thioredoxin-linked reductive inactivation of venom neurotoxins. Mar 1994

11/6/48 08093536 93292511 PMID: 8513799
 Comparison of crotoxin isoforms reveals that stability of the complex plays a major role in its pharmacological action. Jun 1 1993

11/6/49 08081260 94066007 PMID: 8246147
 Chelation of zinc antagonizes the neuromuscular blocking properties of the seven serotypes of botulinum neurotoxin as well as tetanus toxin. Nov 1993

11/6/50 08079555 94024928 PMID: 8212031
 An investigation into the antigenic cross-reactivity of *Ophiophagus hannah* (king cobra) venom neurotoxin, phospholipase A2, hemorrhagin and L-amino acid oxidase using enzyme-linked immunosorbent assay. Jul 1993

11/6/51 07896624 93266553 PMID: 8496167
 Botulinum toxin inhibits arachidonic acid release associated with acetylcholine release from PC12 cells. May 25 1993

11/6/52 07878717 93233028 PMID: 8474024
 Induction of tolerance to crotoxin in mice. Apr 1993

11/6/53 07857082 93190313 PMID: 8446959
 Identification of the site at which phospholipase A2 neurotoxins localize to produce their neuromuscular blocking effects. Jan 1993

11/6/54 07838119 92055781 PMID: 1949068
 Specific binding of crotoxin to brain synaptosomes and synaptosomal membranes. 1991

11/6/55 07837715 92043716 PMID: 1719078
 Role of pertussis toxin-sensitive G proteins in stimulus-dependent human eosinophil degranulation. Nov 15 1991

11/6/56 07833071 91285382 PMID: 2060767
 Properties of a protease-sensitive acceptor component in mouse brain synaptosomes for *Clostridium botulinum* type B neurotoxin. Apr 15 1991

- 11/6/57 07806578 92375369 PMID: 1508413
Ca(2+)-dependent processes as mediators of neurotoxicity. Spring 1992
- 11/6/58 07749556 92342153 PMID: 1378931
Immunochemical analysis of a snake venom phospholipase A2 neurotoxin, crotoxin, with monoclonal antibodies. Jul-Aug 1992
- 11/6/59 07684389 93134619 PMID: 1485347
An EDTA-Ca2+ complex inhibits the enzymatic activity but not the lethality of beta-bungarotoxin. Nov 1992
- 11/6/60 07558975 92085076 PMID: 1727438
Differential effects of presynaptic phospholipase A2 neurotoxins on Torpedo synaptosomes. Jan 1992
- 11/6/61 07529229 91297678 PMID: 1648801
Comparative effects of phospholipase A2 neurotoxins and enzymes on membrane potential and Na+/K+ ATPase activity in rat brain synaptosomes. Jun 15 1991
- 11/6/62 07485424 92104403 PMID: 1761165
Crotoxin, a phospholipase A2 neurotoxin from snake venom, interacts with epithelial mammary cells, is internalized and induces secretion. Nov 1991
- 11/6/63 07443481 91359128 PMID: 1886031
[Study on eosinophil cationic protein (ECP) and arylsulfatase B in nasal secretions and sera from patients with nasal allergy] Jun 1991
- 11/6/64 07424542 91335003 PMID: 1871435
[Eosinophil-induced lung injury] Jun 1991
- 11/6/65 07421762 91329384 PMID: 1868083
Multiplicity of acidic subunit isoforms of crotoxin, the phospholipase A2 neurotoxin from *Crotalus durissus terrificus* venom, results from posttranslational modifications. Aug 13 1991
- 11/6/66 07417132 91320357 PMID: 1862520
Immunochemical cross-reactivity of two phospholipase A2 neurotoxins, agkistrodotoxin and crotoxin. 1991
- 11/6/67 07390190 91198145 PMID: 2015302
Analysis of cDNAs encoding the two subunits of crotoxin, a phospholipase A2 neurotoxin from rattlesnake venom: the acidic non enzymatic subunit derives from a phospholipase A2-like precursor. Mar 26 1991
- 11/6/68 07357812 91050835 PMID: 2146857
Purification and characterization of a phospholipase A2 from human osteoarthritic synovial fluid. 1990
- 11/6/69 07345356 90213209 PMID: 2157526
Inhibition of phosphoinositide hydrolysis by the novel neurotoxin beta-N-oxalyl-L-alpha, beta-diaminopropionic acid (L-BOAA). Feb 26 1990
- 11/6/70 07299955 91089229 PMID: 2264068
In vivo interactions between neurotoxin, cardiotoxin and phospholipases A2 isolated from Malayan cobra (*Naja naja sputatrix*) venom. 1990
- 11/6/71 07237651 90344823 PMID: 2383565
A scorpion venom neurotoxin paralytic to insects that affects sodium current inactivation: purification, primary structure, and mode of action. Jun 26 1990
- 11/6/72 07220030 90225798 PMID: 2327788
The amino acid sequence of a myotoxic phospholipase from the venom of *Bothrops asper*. May 1 1990
- 11/6/73 07161017 91104864 PMID: 2176853
Beta-bungarotoxin-mediated liposome fusion: spectroscopic characterization by fluorescence and ESR. Oct 16 1990
- 11/6/74 07082095 94002063 PMID: 8399216
Model for the interaction of crotoxin, a phospholipase A2 neurotoxin, with presynaptic membranes. Oct 12 1993
- 11/6/75 07034389 93160241 PMID: 8431471
Studies on the subunit structure of textilotoxin, a potent presynaptic neurotoxin from the venom of the Australian common brown snake (*Pseudonaja textilis*). 3. The complete amino-acid sequences of all the subunits. Feb 13 1993
- 11/6/76 07029531 93145974 PMID: 8425551
Snake-venom phospholipase A2 neurotoxins. Potentiation of a single-chain neurotoxin by the chaperon subunit of a two-component neurotoxin. Jan 15 1993
- 11/6/77 07019320 93134601 PMID: 1485334
Isolation and physiological characterization of taicatoxin, a complex toxin with specific effects on calcium channels. Nov 1992
- 11/6/78 07016660 93057502 PMID: 1431893
Effect of phosphorylation of synapsin I and other synaptosomal proteins by beta-bungarotoxin, a phospholipase A2 neurotoxin. Dec 1992
- 11/6/79 06982063 90181446 PMID: 2310754
The complete sequence of the acidic subunit from Mojave toxin determined by Edman degradation and mass spectrometry. Mar 1 1990
- 11/6/80 06938776 94108396 PMID: 1342601
The effect of crotoxin on the release of acetylcholine and lactate dehydrogenase from rat brain cortical slices. 1992
- 11/6/81 06875290 92390948 PMID: 1519253
Specificity of action of beta-bungarotoxin on acetylcholine release from synaptosomes. May-Jun 1992
- 11/6/82 06757645 91263101 PMID: 2048140
Characterization and amino acid sequences of two lethal peptides isolated from venom of Wagler's pit viper, *Trimeresurus wagleri*. 1991
- 11/6/83 06720152 91066875 PMID: 2174508
Mode of action of alpha-latrotoxin: role of divalent cations in Ca2(+)-dependent and Ca2(+)-independent effects mediated by the toxin. Dec 1990
- 11/6/84 06482058 88305766 PMID: 3406946
Mojave toxin: rapid purification, heterogeneity and resistance to denaturation by urea. 1988
- 11/6/85 06319389 87293010 PMID: 3617077
Characterization of the structure and function of three phospholipases A2 from the venom of *Agkistrodon halys pallas*. 1987
- 11/6/86 06310700 87271131 PMID: 3606821
Sequence homology between phospholipase and its inhibitor in snake venom. The primary structure of phospholipase A2 of vipoxin from the venom of the Bulgarian viper (*Vipera ammodytes ammodytes*, Serpentes). Apr 1987
- 11/6/87 06273264 87100994 PMID: 3801413
Characterization of elapidae snake venom components using optimized reverse-phase high-performance liquid chromatographic conditions and screening assays for alpha-neurotoxin and phospholipase A2 activities. Nov 4 1986
- 11/6/88 06073289 89361453 PMID: 2769265
Binding of crotoxin, a presynaptic phospholipase A2 neurotoxin, to negatively charged phospholipid vesicles. Oct 1989
- 11/6/89 05901254 87271931 PMID: 3607213
Interpretation of fluorescence decays in proteins using continuous lifetime distributions. Jun 1987
- 11/6/90 05761701 87250531 PMID: 3036839
The interaction between the presynaptic phospholipase neurotoxins beta-bungarotoxin and crotoxin and mixed detergent-phosphatidylcholine micelles. A comparison with non-neurotoxic snake venom phospholipases A2. Jul 5 1987
- 11/6/91 05689180 86306814 PMID: 2943872
Uncoupling of cholinergic synaptic vesicles by the presynaptic toxin beta-bungarotoxin. Oct 1986
- 11/6/92 05574325 89171252 PMID: 2466692
Neutralization of lethal potency and inhibition of enzymatic activity of a phospholipase A2 neurotoxin, crotoxin, by non-precipitating antibodies (Fab). Feb 13 1989
- 11/6/93 05541946 88025221 PMID: 2444266
[Effect of melittin on ion currents in heart cell membranes] Daistvie melitina na ionnye toki membran kletok serdtsa. Jul-Aug 1987
- 11/6/94 05524017 92206154 PMID: 2562459
Crotoxin, half-century of investigations on a phospholipase A2 neurotoxin. 1989
- 11/6/95 05392446 89255161 PMID: 2656663
Amino acid sequence of a presynaptic neurotoxin, agkistrodotoxin, from the venom of *Agkistrodon halys pallas*. Feb 1989
- 11/6/96 05354471 90089371 PMID: 2597666
Binding of divalent and trivalent cations with crotoxin and with its phospholipase and its non-catalytic subunits: effects on enzymatic activity and on the interaction of phospholipase component with phospholipids. Nov 28 1989
- 11/6/97 05331718 90037075 PMID: 2808427
Beta-bungarotoxin. Preparation and characterization of crystals suitable for structural analysis. Nov 15 1989
- 11/6/98 05324569 90020298 PMID: 2799834
Preparation and characterization of monoclonal antibodies against beta-bungarotoxin and its A- and B-chains. 1989
- 11/6/58 05281957 89318243 PMID: 2749757
Neurotoxic and non-toxic effects of a toxic phospholipase A2 and its nontoxic homologue from the venom of the sea snake, *Laticauda colubrina*. 1989
- 11/6/100 05275806 89302153 PMID: 2742606
[Amino acid sequence of orientotoxins I and II from the venom of the hornet *Vespa orientalis*] Aminokislotaia posledovatel'nost' orientotoksinov I i II iz iada shershnia *Vespa orientalis*. Jan 1989
- 11/6/101 05187376 88163660 PMID: 3349062
Crotoxin, a phospholipase A2 neurotoxin from the South American rattlesnake *Crotalus durissus terrificus*: purification of several isoforms and comparison of their molecular structure and of their biological activities. Jan 26 1988
- 11/6/102 05177701 88107807 PMID: 3337815
Phospholipase A2 hydrolysis of membrane phospholipids causes structural alteration of the nicotinic acetylcholine receptor. Feb 8 1988
- 11/6/103 05110805 88000715 PMID: 3651474
Studies on the subunit structure of textilotoxin, a potent neurotoxin from the venom of the Australian common brown snake (*Pseudonaja textilis*). Sep 24 1987
- 11/6/104 05067511 87184609 PMID: 3566763
Binding ability of *Clostridium botulinum* neurotoxin to the synaptosome upon treatment of various kinds of the enzymes. Mar 30 1987
- 11/6/105 04973884 86104201 PMID: 4084559
Rattlesnake presynaptic neurotoxins: primary structure and evolutionary origin of the acidic subunit. Dec 3 1985
- 11/6/106 04827133 83039480 PMID: 6182570
Antibodies to beta-bungarotoxin and its phospholipase inactive derivative. Jun 22 1982
- 11/6/107 04821180 84261522 PMID: 6204870
Characterization of monoclonal antibodies against beta-bungarotoxin and their use as structural probes for related phospholipase A2 enzymes and presynaptic phospholipase neurotoxins. Jul 2 1984
- 11/6/108 04577164 85028863 PMID: 6489936
Sequence homology between phospholipase and its inhibitor in snake venom. The primary structure of the inhibitor of vipoxin from the venom of the Bulgarian viper (*Vipera ammodytes ammodytes*, Serpentes). Aug 1984
- 11/6/109 04527257 84074091 PMID: 6648985
Ceruleotoxin: identification in the venom of *Bungarus fasciatus*, molecular properties and importance of phospholipase A2 activity for neurotoxicity. 1983
- 11/6/110 04469677 83132074 PMID: 6825753
A comparison of nerve transection and chronic application of beta-bungarotoxin on acetylcholine receptor distribution and other nerve-muscle properties. Mar 1983
- 11/6/111 04430400 82261658 PMID: 7049694
Amino acid sequence of the alpha-subunit of taipoxin, an extremely potent presynaptic neurotoxin from the Australian snake taipan (*Oxyuranus s. scutellatus*). Jun 1982
- 11/6/112 04337331 83079311 PMID: 7173209
Preparation of neurotoxic 3H-beta-bungarotoxin: demonstration of saturable binding to brain synapses and its inhibition by toxin I. Nov 1982
- 11/6/113 04282438 82200155 PMID: 7080027
Isolation of "ceruleotoxin" from *Bungarus fasciatus* venoms. 1982
- 11/6/114 04282437 82200154 PMID: 7080026
Synergism of the two subunits of crotoxin. 1982
- 11/6/115 04202883 81168081 PMID: 7217037
Amino acid sequence of phospholipase A from *Bungarus multicinctus* venom. Jan 1981
- 11/6/116 04062477 83076333 PMID: 7171991
Beta-bungarotoxin-induced cell-death of neurons in chick retina. Nov 4 1982
- 11/6/117 03822457 84104993 PMID: 6661452
[Release of 14C-acetylcholine from synaptosomes as affected by presynaptic neurotoxins—from bee and cobra venoms—phospholipases A2] Vysvobozhdenie [14C] atsetilkholina iz sinaptosom pod vlianiem presinapticheskikh neirotoksinov iz iadov zmei i pchely—fosfolipaz A2. Nov 1983
- 11/6/118 03560161 80175361 PMID: 7370229
Phospholipase A2 activity and substrate specificity of snake venom presynaptic toxins. Mar 18 1980
- 11/6/119 03545041 80109855 PMID: 7353053
Penetrability of proteins through the digestive system of *Sarcophaga falculata* blowfly. Jan 3 1980

- 11/6/120 03385405 76102740 PMID: 1749
beta-Bungarotoxin, a pre-synaptic toxin with enzymatic activity. Jan 1976
- 11/6/121 03176286 79222410 PMID: 460683
Neuronal degeneration induced by stereotaxic injection of beta-bungarotoxin into rat brain. Feb 1979
- 11/6/122 03017806 77087738 PMID: 831773
Relationship between the neurotoxicity and phospholipase A activity of beta-bungarotoxin. Jan 11 1977
- 11/6/123 02897111 76206154 PMID: 1276197
Saturable binding to cell membranes of the presynaptic neurotoxin, beta-bungarotoxin. May 21 1976
- 11/6/124 02869581 76092395 PMID: 1245225
The role of phospholipase activity in the action of a presynaptic neurotoxin from the venom of *Notechis scutatus scutatus* (Australian tiger snake). Jan 1 1976
- 11/7/46 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
08140009 94223544 PMID: 8169833
Inhibition of vacuolar adenosine triphosphatase antagonizes the effects of clostridial neurotoxins but not phospholipase A2 neurotoxins.
Simpson LL; Coffield JA; Bakry N
Department of Medicine, Jefferson Medical College, Philadelphia, Pennsylvania.
Journal of pharmacology and experimental therapeutics (UNITED STATES) Apr 1994, 269 (1) p256-62, ISSN 0022-3565 Journal Code: JP3 Contract/Grant No.: NS-22153, NS, NINDS Languages: ENGLISH Document type: Journal Article Record type: Completed
Bafilomycin A1, an inhibitor of vacuolar adenosine triphosphatase, was tested for its ability to antagonize botulinum neurotoxins (serotypes A-G), tetanus toxin and phospholipase A2 neurotoxins (notexin, beta-bungarotoxin, taipoxin and textilotoxin) on the mouse phrenic nerve-hemidiaphragm preparation. Bafilomycin itself produced concentration-dependent blockade of neuromuscular transmission without blocking nerve action potentials or muscle action potentials. This effect may have been due to inhibition of the proton pump that regulates acetylcholine transport into vesicles. At submaximal concentrations, bafilomycin was very effective in delaying the onset of paralysis due to all clostridial neurotoxins, but it had no protective effect against phospholipase A2 neurotoxins. Experiments were done to determine which of the three steps in clostridial neurotoxin action was antagonized by bafilomycin (e.g., binding, internalization and intracellular poisoning). Both pharmacological experiments and ligand-binding experiments showed that the drug did not block toxin binding to the plasma membrane. Similarly, pharmacological experiments on the time-dependent effects of bafilomycin showed that the drug did not antagonize the intracellular actions of toxins. The data indicated that bafilomycin acted at the intermediate step of internalization. This is in keeping with the facts that: 1) bafilomycin inhibits vacuolar adenosine triphosphatase, which in turn leads to inhibition of acidification in endosomes and 2) clostridial neurotoxins depend upon acidification of endosomes for translocation to the cytosol. The finding that bafilomycin antagonizes tetanus toxin may provide important clues for understanding how this toxin can act locally to produce flaccid paralysis. The finding that bafilomycin is a universal antagonist that protects against all clostridial neurotoxins may have important implications for developing therapeutic drugs. Record Date Created: 19940602
- 11/5/49 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
08081260 94066007 PMID: 8246147
Chelation of zinc antagonizes the neuromuscular blocking properties of the seven serotypes of botulinum neurotoxin as well as tetanus toxin.
Simpson LL; Coffield JA; Bakry N
Department of Medicine, Jefferson Medical College, Philadelphia, Pennsylvania.
Journal of pharmacology and experimental therapeutics (UNITED STATES) Nov 1993, 267 (2) p720-7, ISSN 0022-3565 Journal Code: JP3 Contract/Grant No.: NINDS NS-22153, NS, NINDS Languages: ENGLISH Document type: Journal Article Record type: Completed
Botulinum neurotoxin types A, B (unactivated and activated), C, D, E, F and G, as well as tetanus toxin, paralyzed transmission in mouse phrenic nerve-hemidiaphragm preparations. Toxin-induced blockade of transmission was antagonized by chelators [e.g., ethylenediamine tetraacetic acid, tetrakis(2-pyridylmethyl)ethylenediamine or diethylene-triaminepentaacetic anhydride], but
- 17/6/1 0817918 20332947 PMID: 10874773
Ocular aspects of myasthenia gravis. 2000
- 17/6/2 10358931 20000569 PMID: 10532769
Strabismus surgery among aged medicare beneficiaries. Dec 1997
- 17/6/3 10240117 99371424 PMID: 10443824
The posterior thyroplasty window: anatomical considerations. Aug 1999
- 17/6/4 09825762 98378683 PMID: 9713061
Botulinum toxin A treatment of overactive corrugator supercilii in thyroid eye disease. May 1998
- 17/6/5 09428059 98018536 PMID: 9380360
Chemodeneration in treatment of upper eyelid retraction. 1997
- 17/6/6 08164191 94255188 PMID: 8196925
Control of eyelid retraction associated with Graves' disease with botulinum A toxin. Mar 1994
- 11/6/125 02826941 76005594 PMID: 1158892
Amino acid sequence of a presynaptic neurotoxin from the venom of *Notechis scutatus scutatus* (Australian tiger snake). Sep 10 1975
- 11/6/126 02758131 77138542 PMID: 191284
Snake venom action: are enzymes involved in it? Feb 15 1977
- 11/6/127 02648802 80043375 PMID: 387107
Effect of presynaptic neurotoxin notechis II-5 from tiger snake venom on the motor nerve endings of mice] Deistvie presinapticheskogo neirotoksina notechisa II-5 iz iada tigrovoi zmei na dvigatel'nye nervnye okonchaniia myshi. Oct 1979
- 11/6/128 02559054 80046716 PMID: 499210
Postsynaptic effects of crotoxin and of its isolated subunits. Sep 1979
- 11/6/129 02557313 80043177 PMID: 497256
Amino acid sequence of a postsynaptic neurotoxin from the venom of the Australian tiger snake *Notechis scutatus scutatus*. 1979
- this effect was dependent on incubation conditions. Pretreatment of toxin with chelators failed to produce antagonism, but pretreatment of tissues did produce antagonism. Of the various chelators tested, tetrakis(2-pyridylmethyl)ethylenediamine produced the greatest effect. Antagonism of toxin-induced neuromuscular blockade could be partially reversed by washing chelators from tissues and could be fully reversed by adding an excess of zinc. The ability of chelators to antagonize clostridial neurotoxins was specific and did not extend to phospholipase A2 neurotoxins. Ligand-binding studies with radioiodinated toxin and brain membrane preparations showed that chelators did not antagonize toxicity by inhibiting toxin association with receptors. Similarly, pharmacological experiments with unlabeled toxin- and type-specific antibodies demonstrated that chelators did not act by blocking receptor-mediated internalization of toxin. The chelators appeared to exert their effects by antagonizing the intracellular actions of clostridial neurotoxins. Electrophysiological studies showed that chelators, at concentrations relevant to antagonism of botulinum neurotoxin and tetanus toxin, did not enhance transmitter release.(ABSTRACT TRUNCATED AT 250 WORDS) Record Date Created: 19940106
Tags: Animal; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.
Descriptors: *Botulinum Toxins-antagonists and inhibitors-AI; *Chelating Agents-pharmacology-PD; *Neuromuscular Blocking Agents-antagonists and inhibitors-AI; *Tetanus Toxin-antagonists and inhibitors-AI; *Zinc-metabolism-ME; Botulinum Toxins-classification-CL; Botulinum Toxins-metabolism-ME; Brain-metabolism-ME; Chelating Agents-metabolism-ME; Metalloendopeptidases-metabolism-ME; Mice; Neuromuscular Junction-drug effects-DE; Neuromuscular Junction-metabolism-ME; Rats; Sensitivity and Specificity; Serotyping; Tetanus Toxin-classification-CL; Tetanus Toxin-metabolism-ME CAS Registry No.: 0 (Botulinum Toxins); 0 (Chelating Agents); 0 (Neuromuscular Blocking Agents); 0 (Tetanus Toxin); 7440-66-6 (Zinc)
Enzyme No.: EC 3.4.24 (Metalloendopeptidases)
- 11/7/104 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
05067511 87184609 PMID: 3566763
Binding ability of Clostridium botulinum neurotoxin to the synaptosome upon treatment of various kinds of the enzymes.
Kitamura M; Sone S
Biochemical and biophysical research communications (UNITED STATES) Mar 30 1987, 143 (3) p928-33, ISSN 0006-291X Journal Code: 9Y8 Languages: ENGLISH Document type: Journal Article Record type: Completed
The binding ability of Cl. botulinum neurotoxin to synaptosomes upon treatment with various enzymes (neuraminidase, trypsin, and beta-bungarotoxin containing phospholipase A2 activity) was studied. When synaptosomes were treated with neuraminidase, their ability to bind toxin decreased; trypsin and beta-bungarotoxin had slightly weak or no effect. The decrease in toxin-binding ability of synaptosomes was paralleled by a release of sialic acid from the synaptosomes by the neuraminidase treatment. The toxin-binding ability of synaptosomes treated with neuraminidase was lower than untreated ones at a high concentration of sodium chloride. The binding of the toxin to synaptosomes occurred at least at the two types of structural sites, one site which contained sialic acid, and other site which was sensitive to high ionic strength. It may be possible that another binding state except these is present at the synapse. Record Date Created: 19870518
- 17/6/13 06008165 86099357 PMID: 3002078
[Myasthenic syndromes linked to mediator secretion disorders] Miastenicheskie sindromy, svyazannyye s narusheniem sekretsii mediatora. 1985
- 17/6/14 05312485 89385497 PMID: 2779991
Botulinum toxin therapy of eye muscle disorders. Safety and effectiveness. American Academy of Ophthalmology. Sep 1989
- 17/6/15 05204825 89026356 PMID: 3179055
Botulinum toxin. Aug 1988
- 17/6/16 05131078 87070940 PMID: 3466462
Diplopia in thyroid eye disease. 1986
- 17/6/17 04905297 84225515 PMID: 6676980
Saccadic velocity measurements in strabismus. 1983
- 17/6/18 04576471 85026678 PMID: 6489104
Injection treatment of endocrine orbital myopathy. Aug 15 1984
- 17/7/3 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.

10240117 99371424 PMID: 10443824

The posterior thyroplasty window: anatomical considerations.

Maragos NE

Mayo Clinic, Rochester, Minnesota 55905, USA.

Laryngoscope (UNITED STATES) Aug 1999, 109 (8) p1228-31, ISSN 0023-852X

Journal Code: L1W Languages: ENGLISH Document type: Journal Article Record type:

Completed

OBJECTIVES: Explain surgical technique of performing a posterior thyroplasty window. Describe the internal laryngeal anatomy and structures available through the posterior window approach. Describe posterior window approach. **STUDY DESIGN:** Review of lateral laryngeal anatomy and retrospective review of 125 cases involving a posterior thyroplasty window approach. Review mechanics of stress and stress concentration inherent with partial removal of rigid substance. Describe anatomical considerations and surgical complications. **METHODS:** Charts were reviewed and tabulated for surgical complications, efficacy and safety of surgical approach, specific anatomical variations, and variety of surgery available through the posterior window. **RESULTS:** Performance of 125 posterior thyroplasty windows revealed no evidence of entry into the piriform sinus. Three thyroid ala fractures ensued, two of the body and one of the inferior cornu. Operations available included arytenoid adduction, arytenoid fixation, lysis of joint adhesions, and access to the posterior cricoarytenoid muscle for botulinum toxin injections. **CONCLUSIONS:** The posterior thyroplasty window affords easy, direct access to the internal, posterolateral larynx while preserving the cricothyroid joint, the action of the cricothyroid muscle, and the internal division of the recurrent laryngeal nerve. Record Date Created: 19990907

1777/8 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
06895291 93064763 PMID: 1437204

Bilateral thyroarytenoid denervation: a new treatment for laryngeal hyperadduction disorders studied in the canine.

Sercarz JA; Berke GS; Ming Y; Rothschilder J; Graves MC

UCLA School of Medicine, Division of Head and Neck Surgery.

Otolaryngology--head and neck surgery (UNITED STATES) Nov 1992, 107 (5) p657-68,

ISSN 0194-5998 Journal Code: ON8 Languages: ENGLISH Document type: Journal Article

Record type: Completed

Adductor spasmodic dysphonia is a vocal disorder of uncertain etiology with no satisfactory long-term treatment. Recently, injection of botulinum toxin (Botox) into the thyroarytenoid (TA) muscle has been used as an effective temporary treatment. A surgical counterpart to bilateral TA Botox injection is described in this article. Bilateral thyroarytenoid denervation was performed through a window in the thyroid cartilage in seven canines, including four that were studied 3 months after the procedure. No serious complications occurred in the animals, each maintaining full vocal fold abduction and adduction. In all cases, anticipated physiologic changes in laryngeal function were observed, including the inability to generate high subglottic pressures during high levels of laryngeal nerve (RLN) stimulation. In two of the surviving animals, the ansa cervicalis was used to reinnervate the TA muscle, thereby preventing the possibility of reinnervation from the proximal RLN stump while limiting TA atrophy and fibrosis. Bilateral TA denervation represents a hopeful new long-term approach to spasmodic dysphonia treatment. Record Date Created: 19921218

1777/15 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
05204825 89026356 PMID: 3179055

Botulinum toxin.

Kowal L

Australian and New Zealand journal of ophthalmology (AUSTRALIA) Aug 1988, 16 (3)

p264-6, ISSN 0814-9763 Journal Code: ANZ Languages: ENGLISH Document type: Journal

Article Record type: Completed Record Date Created: 19881220

1777/16 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv.
05131078 87070940 PMID: 3466462

Diplopia in thyroid eye disease.

Fells P; McCarty B

Transactions of the ophthalmological societies of the United Kingdom (ENGLAND) 1986,

105 (Pt 4) p413-23, ISSN 0078-5334 Journal Code: WA1 Languages: ENGLISH Document

type: Journal Article Record type: Completed Record Date Created: 19870122

S9 23407 NEUROTOX? S10 0 S7 AND S9 S11 1540602 TREAT? S12 321 S11 AND S7

02Jul01 08:32:48 User208600 Session D1405.1 File 155:MEDLINE(R) 1966-2001Jul W2 (c) format only 2001 Dialog Corporation

Set Items Description S1 6430 BOTUL? S2 17401 CALCITONIN S3 8 S1 AND S2 S4 19552 HYPOCALC? OR HYPERCALC? S5 5 S1 AND S4 S6 90754 THYROID S7 1231 S4 AND S6 S8 0 S1 AND S7

3/6/1 09948794 98433827 PMID: 9762863 Effects of botulinum neurotoxin type A on the expression of gephyrin in cat abducens motoneurons. Oct 12 1998

3/6/2 09471682 98014520 PMID: 9344569 Sensory and motor denervation influence epidermal thickness in rat foot glabrous skin. Oct 1997

3/6/3 09404508 97291189 PMID: 9145803 Expression of neurotransmitter genes in rat spinal motoneurons after chemodenervation with botulinum toxin. May 1997

3/6/4 09159891 97110858 PMID: 9081635

3/7/5 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 09088502 96364144 PMID: 8739382

Upregulation of calcitonin gene-related peptide at mouse motor nerve terminals poisoned with botulinum type-A toxin. Meunier FA; Colasante C; Faillie L; Gastard M; Molgo J Laboratoire de Neurobiologie Cellulaire et Moléculaire, C.N.R.S. (U.P.R.9040), Gif-sur-Yvette, France. Pflugers Archiv (GERMANY) 1996, 431 (6 Suppl 2), pR297-8, ISSN 0031-6768 Journal Code: OZX Languages: ENGLISH Document type: Journal Article Record type: Completed

Calcitonin gene-related peptide (CGRP)-like immunoreactivity of motor nerve terminals was investigated at different times after local in vivo injection of botulinum type-A toxin (BoNT/A) close to the mouse levator auris longus muscle. CGRP expression in most of control nerve terminals was undetectable, but markedly increased during muscle paralysis and synaptic remodelling and declined once functional recovery occurred. Record Date Created: 19970116

5/7/1 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 10922488 20471417 PMID: 11021434

Fatal Clostridium botulinum toxicosis in eleven Holstein cattle fed round bale barley haylage. Kelch WJ; Kerr LA; Pingle JK; Rohrbach BW; Whitlock RH Department of Comparative Medicine, College of Veterinary Medicine, University of Tennessee, Knoxville 37901-1071, USA. Journal of veterinary diagnostic investigation (UNITED STATES) Sep 2000, 12 (5) p453-5, ISSN 1040-6387 Journal Code: AZD Languages: ENGLISH Document type: Journal Article Record type: Completed

Twenty-two lactating Holstein cattle in Tennessee had clinical signs of intoxication with preformed Clostridium botulinum toxin. These signs included weakness, paralysis of the tongue and chest muscles, abdominal breathing, and, in 11 of the 22 cows, death. Differential diagnoses included hypocalcemia, hypomagnesemia, carbohydrate overload, and several toxicooses including mycotoxin, lead, nitrate, organophosphate, atropine or atropine-like alkaloid, and botulism. A diagnosis of botulism by the ingestion of preformed C. botulinum type B toxin was made by eliminating these other diseases, by finding C. botulinum type B spores in 3 bales of round bale barley haylage fed to these cattle, and by isolating preformed type B toxin from 1 of the 3 bales. Confirmation of the toxin type was made by demonstrating mouse lethality by intraperitoneal injection of specimen extracts with neutralization by C. botulinum type B antitoxin. The haylage, harvested green and encased in black plastic bags to facilitate fermentation, was presumably contaminated by the botulinum toxin when fermentation failed to produce enough acid to lower the pH to 4.5, the pH below which C. botulinum growth is inhibited. Farmers and ranchers who use round hay balers to produce haylage should be alert to this potential problem. Record Date Created: 20010108

5/7/2 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 08679305 96124965 PMID: 8526819

The clinical differentiation of nervous and muscular locomotor disorders of sheep in Australia. Bourke CA 5/6/4 06402894 87186703 PMID: 3494493

Localized diseases of the bovine brain and spinal cord. Mar 1987

5/6/5 03657791 83094030 PMID: 7348919 [Central nervous system diseases in cattle. 2. Diseases in young and adult cattle] Erkrankungen des Zentralnervensystems beim Rind. 2. Die Krankheiten jugendlicher und erwachsener Rinder. 1981

12/6/1 11321719 21280324 PMID: 11386024 Coexistence of parathyroid carcinoma and non-medullary carcinoma of the thyroid. May-Jun 2001

12/6/2 11371761 21270595 PMID: 11376402 Minima ly invasive video-assisted parathyroidectomy: multinstitutional study. Jun 2001

12/6/3 11303912 21129989 PMID: 11234312 Short- and long-term results of total vs subtotal thyroidectomies in the surgical treatment of Graves' disease. 2001

12/6/4 11303377 21132265 PMID: 11236346 Incidence of complications of thyroid surgery [Incidence delle complicanze nella chirurgia della tiroide. Oct 2000

12/6/5 11256203 21241882 PMID: 11344634

Regulation of motoneuronal calcitonin gene-related peptide (CGRP) during axonal growth and neuromuscular synaptic plasticity induced by botulinum toxin in rats. Apr 1996

3/6/5 09088502 96364144 PMID: 8739382 Upregulation of calcitonin gene-related peptide at mouse motor nerve terminals poisoned with botulinum type-A toxin. 1996

3/6/6 08461157 95123477 PMID: 7623160 Calcitonin gene-related peptide: possible role in formation and maintenance of neuromuscular junctions. Jan 1995

3/6/7 08356509 95187010 PMID: 7881294 Calcitonin gene-related peptide-like immunoreactivity, in botulinum toxin-paralysed rat muscles. Sep-Nov 1994

3/6/8 07960661 94061378 PMID: 8242236 Enhancement by calcitonin gene-related peptide of nicotinic receptor-operated noncontractile Ca2+ mobilization at the mouse neuromuscular junction. Oct 1993

NSW Agriculture, Agricultural Research and Veterinary Centre. Australian veterinary journal (AUSTRALIA) Jun 1995, 72 (6) p228-34, ISSN 0005-0423 Journal Code: 9IE Languages: ENGLISH Document type: Journal Article; Review, Tutorial Record type: Completed

Many of the nervous and muscular locomotor disorders that affect sheep throughout Australia are commonly referred to as "stagers" syndromes. The range of clinical signs displayed by sheep suffering these disorders is sufficiently diverse to enable each syndrome to be graded into one of 5 progressive clinical groups. The first group, the limb paresis syndromes, includes the primary myopathies associated with the ingestion of Ixobrya brevicornis, Malva parviflora, and Trachymene ochracea, as well as selenium and Vitamin E disorders, Paro virus staggers, congenital progressive muscular dystrophy, humpy back, hypocalcaemic muscle weakness, Tribulus terrestris staggers and tetanus. The second group is characterised by limb paresis with knuckling of the fetlocks, and includes the plant-associated toxicities of Romulea rosea, Stachys arvensis, Trachyantha divaricata, and Tribulus micrococcus, together with haloxon toxicity, enzootic ataxia (copper deficiency), and the probably genetic disorders of segmental axonopathy, neuroaxonal dystrophy, and degenerative thoracic myelopathy. Other locomotor disorders that fit more loosely into this group are islet myelitis (post-dipping staggers), vitamin A deficiency, cervico-thoracic vertebral subluxation Stypandra glauca toxicity, Ipomoea spp toxicity, ivermectin toxicity, and botulism. The third group, the falling syndromes, includes the probably genetic disorders of thalamic cerebellar neuropathy, cerebellar atrophy, and globoid cell leucodystrophy, together with Swainsona spp toxicity, the fourth group, the falling syndromes, includes the plant associated toxicities of phalaris staggers, perennial ryegrass staggers and nervous ergotism (Claviceps paspali). (ABSTRACT TRUNCATED AT 250 WORDS) (40 Refs.) Record Date Created: 19960122

5/7/3 DIALOG(R)File 155:MEDLINE(R) (c) format only 2001 Dialog Corporation. All rts. reserv. 07689557 93149376 PMID: 1491765

[Primary hyperparathyroidism with prevalent neuro-muscular manifestations] [iparatiroidismo primario con prevalenti manifestazioni neuro-muscolari. Bartolucci L; Fioretti M; Fiorini E; Proietti MG; Gradoli C; Valoni C Istituto di Patologia Speciale Medica, Università degli Studi di Perugia, Ospedale Civile, Terni. Minerva medica (ITALY) Dec 1992, 83 (12) p841-5, ISSN 0026-4806 Journal Code: N6M Languages: ITALIAN Document type: Journal Article Record type: Completed

A case of primary hyperparathyroidism with prevalent neuromuscular symptoms is described. Clinical, diagnostic and therapeutic implications are emphasized. Particular attention must involve a full clinical examination, electromyographic da neuromuscular biopsy to make differentiation from primary myopathy or denervation pathology. Some similarity of electromyographic data with those observed in botulism and myasthenia gravis should also be taken in mind. Hypercalcaemia could play a pathological role in conditioning abnormalities of nervous impulse conduction at the level of neuromuscular junction. Another possible interference might be related to a direct effect of parathormone and hypophosphataemia on nervous impulse conduction. "Glandular hyperplasia", as observed in this case at istologic examination, rises some problems as far as the prognosis is concerned. Record Date Created: 19930304

[In Process Citation] Akutni pankreatitida jako cesta k diagnoze primarni hyperparatyreoz. Oct 2000

12/6/6 11232524 21226317 PMID: 11328475 Review of the outcome of management of Graves' disease in children and adolescents. Apr 2001

12/6/7 11229438 21223268 PMID: 11323564 Parathyroid carcinoma. Therapeutic strategies derived from 20 years of experience. Mar 2001

12/6/8 11229226 21222928 PMID: 11323743

Extracellular calcium-sensing receptor: structural and functional features and association with diseases. May 2001

126/9 11210613 21213281 PMID: 11315456
Endocrine problems in the chronically critically ill patient. Mar 2001

126/10 11132017 21071012 PMID: 11190548
Post-thyroidectomy hypocalcaemia and feasibility of short-stay thyroid surgery. Sep-Oct 2000

126/11 11081429 21041817 PMID: 11201007
Primary hyperparathyroidism—early diagnosis in patients referred for thyroid surgery. Dec 2000

126/12 11081275 21094397 PMID: 11177016
Thyroidectomy for selected patients with thyrotoxicosis. Jan 2001

126/13 11070816 21129969 PMID: 11234312
Short- and long-term results of total vs subtotal thyroidectomies in the surgical treatment of Graves' disease. 2001

126/14 11069718 21132265 PMID: 11236346
[In Proc ess Citation] Incidenza delle complicanze nella chirurgia della tiroide. Oct 2000

126/15 11007036 21071012 PMID: 11190548
Post-thyroidectomy hypocalcaemia and feasibility of short-stay thyroid surgery. Sep-Oct 2000

126/16 10958887 21013572 PMID: 11128714
Treatment of a thyrotoxinoma with octreotide-LAR in a patient with multiple endocrine neoplasia-1. Nov 2000

126/17 10911191 21011899 PMID: 11125809
Minimally invasive video-assisted parathyroidectomy: lesson learned from 137 cases. Dec 2000

126/18 10907686 20529367 PMID: 11077587
Standardized morphology-adjusted resection in treatment of benign nodular struma [Die standardisierte morphologiegerechte Resektion in der Behandlung von benignen Knotenstrumen Oct 2000

126/19 10841496 20471218 PMID: 11019711
Hypercalcaemia following renal transplantation in a cat. Oct 1 2000

126/20 10828272 20490451 PMID: 11034808
Clinical aspects of early and late hypocalcaemia after thyroid surgery. Sep 2000

126/21 10818304 20409496 PMID: 10953376
Surgical treatment of rare anatomic variations and anomalies of the thyroid gland [Hirurgsko lecenje relikh anatomskih varijacija i anomalija stitaste zlezde. 2000

126/22 10800626 99157851 PMID: 10048291
Unaccountable severe hypercalcaemia in a patient treated for hypoparathyroidism with dihydroxycholesterol. Jan 1999

126/23 10799327 20376594 PMID: 10921180
[Subsistmal goiter. Report of 212 cases] Les goitres plongeants. A propos de 212 cas. Jan 2000

126/24 10798099 20367017 PMID: 10909288
[Parathyroid neoplasm—diagnostic challenge and therapeutic difficulty] Rak przytarczyc—wyzwanie diagnostyczne i problem terapeutyczny. 2000

126/25 10771726 20036958 PMID: 10567671
Antineoplastic activity of 1,25(OH)2D3 and its analogue 22-oxacalcitriol against human anaplastic thyroid carcinoma cell lines in vitro. Dec 1999

126/26 10748602 98253676 PMID: 9591451
Surgical treatment of secondary hyperparathyroidism in patients treated with renal replacement [Leczenie operacyjne wtornej nadczynności przytarczyc u chorych leczonych nerkozastępczo. 1997

126/27 10729189 20369258 PMID: 10911605
[Intraoperative neurophysiological monitoring (IONM) of the recurrent laryngeal nerve and microdissection. Surgical techniques for decreasing the risk of recurrent laryngeal nerve paralysis] Intraoperative neurophysiologische Monitoring (IONM) des Nervus recurrens

und Mikrodissektion. Operationstechnische Verfahren zur Risikominimierung von Recurrensparesen. May 2000

126/28 10665587 20357060 PMID: 10897019
Initial results from a prospective cohort study of 5583 cases of thyroid carcinoma treated in the united states during 1996. U.S. and German Thyroid Cancer Study Group. An American College of Surgeons Commission on Cancer Patient Care Evaluation study. Jul 1 2000

126/29 10644895 20288129 PMID: 10829320
Effect of preoperative preoperative visualization therapy on postoperative outcome after inguinal hernia surgery and thyroid resection [Einfluss einer präoperativen Visualisierungstherapie auf die postoperativen Ergebnisse nach Leistenherniotomie und Strumaresektion. 2000

126/30 10621282 20272344 PMID: 10812464
Change of surgical treatment in benign thyroid gland diseases— influence of surgical management on perioperative complications [Wandel der chirurgischen Therapie bei benignen Schilddrüsenerkrankungen— Einfluss der chirurgischen Taktik auf perioperative Komplikationen. 2000

126/31 10613077 20237873 PMID: 10775283
Causes of anorexia in untreated hyperthyroidism: a prospective study. May 2000

126/32 10598687 20260383 PMID: 10800589
[A case of pseudohypoparathyroidism] Przypadek rzekomej niedoczynności przytarczyc. 1999

126/33 10531979 20164362 PMID: 10699905
Medullary thyroid carcinoma: clinical characteristics, treatment, prognostic factors, and a comparison of staging systems. Mar 1 2000

126/34 10525997 20182965 PMID: 10720046
Changes in parameters of bone and mineral metabolism during therapy for hyperthyroidism. Mar 2000

126/35 10520327 20159709 PMID: 10695386
[Retrosental parathyroid gland cystic neoplasm as a cause of primary hyperparathyroidism] Rak zamosiłkowej torbieli przytarczyc jako przyczyna pierwotnej nadczynności przytarczyc. 1999

126/36 10506060 20086675 PMID: 10619879
Postoperative complications of surgery for malignant tumors of the thyroid gland [Complicaciones tras la cirugía de los tumores malignos de la glándula tiroidea. Oct 1999

126/37 10501692 20122825 PMID: 10658364
A comparative analysis of the postoperative complications of thyroid cancer surgery related to surgical approach. 1999

126/38 10482262 20119769 PMID: 10654273
Hyperthyroidism after surgery for primary hyperparathyroidism. Dec 1999

126/39 10457106 20082640 PMID: 10615778
[Video-assisted parathyroidectomy: a series of 85 cases] Parathyroidectomy video-assisted: une serie de 85 cas. Nov 1999

126/40 10441923 20001341 PMID: 10633774
Surgical treatment of the substernal goitre. Jul-Sep 1999

126/41 10429748 20066127 PMID: 10598180
Unilateral neck exploration under local anesthesia: the approach of choice for asymptomatic primary hyperparathyroidism. Dec 1999

126/42 10388794 20043094 PMID: 10575840
De novo RET proto-oncogene mutation in a patient with multiple endocrine neoplasia type 2B. Oct 1999

126/43 10368276 99443262 PMID: 10515239
Hypercalcaemia -induced renal insufficiency during therapy with dihydroxycholesterol. 1999

126/44 10285646 99409767 PMID: 10482041
Coincidence of hot thyroid nodules and primary hyperparathyroidism. 1999

126/45 10260975 99387764 PMID: 10460008
Effect of 22-oxa-1,25-dihydroxyvitamin D3 on human thyroid cancer cell growth. Apr 1999

126/46 10253026 99357134 PMID: 10430065
A Phase I study of LGD1069 in adults with advanced cancer. Jul 1999

126/47 10237517 99241664 PMID: 10225153
Parathyroid and thyroid function five years after treatment of laryngeal and hypopharyngeal carcinoma. Apr 1999

126/48
10184010 99295580 PMID: 10367504
Subsistmal goiter: a diagnostic and therapeutic problem. (Report of 39 surgically treated cases) [Gozzo immenso: un problema diagnostico e terapeutico. (Osservazioni su 39 operati). Jan-Feb 1999

126/49 10171189 99283054 PMID: 10354657
Effect of lithium maintenance therapy on thyroid and parathyroid function. May 1999

126/50 10072238 99186032 PMID: 10086113
[Prophylactic total thyroidectomy in childhood for multiple endocrine neoplasia type 2A: preliminary results] Profylaktische totale Thyroidectomie voor multiple endocriene neoplasie type 2A op de kindereeftijd: eerste ervaringen. Jan 9 1999

126/51 09987751 99006143 PMID: 9789714
Effects of parathyroid hormone infusion on glucose tolerance and glucose-stimulated insulin secretion in normal and uremic rats. Aug 1998

126/52 09983746 99071738 PMID: 9654574
Primary and reoperative parathyroid operations in hyperparathyroidism of multiple endocrine neoplasia type 1. Dec 1998

126/53 09964058 99006021 PMID: 9789592
[Morbidity from total thyroidectomy in subjects less than 15 years of age. A multicenter study of 107 cases] Morbidite de la thyroidectomie totale chez le sujet de moins de 15 ans. Etude multicentrique de 107 cas. Jul 1998

126/54 09960466 99040738 PMID: 9823404
Early relapse after operation for Graves' disease: postoperative hormone kinetics and outcome after subtotal, near-total, and total thyroidectomy. Nov 1998

126/55 09927516 98448322 PMID: 9775196
[Aetiological features of osteoporosis in male patients aged less than 50 years: study of 28 cases with a comparative series of 30 patients over the age of 50] Aspects étiologiques de l'ostéoporose masculine avant 50 ans: étude de 28 cas comparés a 30 cas ages de plus de 50 ans. Jul 1998

126/56 09915231 98415705 PMID: 9744456
Parathyroidectomy via bilateral cervical exploration: a retrospective review of 866 cases. Oct 1998

126/57 09906312 98406309 PMID: 9735129
Medullary thyroid carcinoma: genetic advances, treatment recommendations, the approach to the patient with persistent hypercalcinaemia. Oct 1998

126/58 09894163 98367299 PMID: 9701977
[Hypoparathyroidism after thyroidectomy. Analysis of a consecutive, recent series] [Hypoparathyroidismo dopo tiroidectomia. Verifica di una serie consecutiva e recente. Apr 1998

126/59 09894162 98367298 PMID: 9701976
Subtotal thyroidectomy or total thyroidectomy in the treatment of benign thyroid disease. Our experience [Tiroidectomia subtotale o tiroidectomia totale nel trattamento della patologia benigna della tiroide. Nostra esperienza. Apr 1998

126/60 09893018 98357482 PMID: 9694109
The impact of initial surgical management on outcome in young patients with differentiated thyroid cancer. Jul 1998

126/61 09885657 98406607 PMID: 9735646
Surgical treatment of thyroid cancer: the Singapore General Hospital experience. Aug 1998

126/62 09799313 98317987 PMID: 9655281
Thyroid reoperations: indications and risks. Jul 1998

- 126/63 09774936 98249659 PMID: 9588060
[Thyroidectomy without drainage] Thyroidectomy sans drainage. 1997
- 126/64 09755886 98244783 PMID: 9585384
Chronic hypervitaminosis D3 determines a decrease in C-cell numbers and calcitonin levels in rats. Feb 1998
- 126/65 09737244 98203131 PMID: 9542028
Total thyroidectomy in iodine-deficient goiter--an effective treatment alternative? Die totale Thyreoidektomie bei Jodmangelstruma--eine sinnvolle Behandlungsalternative? 1998
- 126/66 09737240 98203127 PMID: 9542024
Rate of complications with systematic exposure of the recurrent laryngeal nerve and parathyroid glands in operations for benign thyroid gland diseases] Die Komplikationsrate bei systematischer Darstellung des Nervus recurrens und der Epithelkörperchen für Operationen benigner Schilddrüsenerkrankungen 1998
- 126/67 09737239 98203124 PMID: 9542021
Surgical therapy of benign thyroid gland diseases] Die chirurgische Therapie benigner Schilddrüsenerkrankungen 1998
- 126/68 09735564 98223006 PMID: 9563542
Differentiated thyroid cancer: determinants of disease progression in patients <21 years of age at diagnosis: a report from the Surgical Discipline Committee of the Children's Cancer Group. Apr 1998
- 126/69 09719051 98166117 PMID: 9505219
Metastatic parathyroid carcinoma (mPCa): natural history and treatment of a case. Dec 1997
- 126/70 09669266 98146317 PMID: 9486896
Safety of completion thyroidectomy for multicentric carcinoma. Feb 1998
- 126/71 09664675 98128178 PMID: 9467125
Carcinoma of the thyroid in children--a review. Nov-Dec 1997
- 126/72 09620131 98072778 PMID: 9408498
Reoperation for missed parathyroid adenoma. 1997
- 126/73 09607039 98072289 PMID: 9409611
Intrathyroidal parathyroid glands can be a cause of failed cervical exploration for hyperparathyroidism. Dec 1997
- 126/74 09601852 98068530 PMID: 9406695
Anti-carcinoembryonic antigen antibodies versus somatostatin analogs in the detection of metastatic medullary thyroid carcinoma: are carcinoembryonic antigen and somatostatin receptor expression prognostic factors? Dec 15 1997
- 126/75 09579381 97442951 PMID: 9297882
[Results of reoperations for goiter] Resultats des reinterventions pour goitres. 1997
- 126/76 09415324 98026749 PMID: 9362426
Skeletal complications of malignancy, Oct 15 1997
- 126/77 09405799 97353588 PMID: 9209829
Amylin, calcitonin gene-related peptide, calcitonin, and adrenomedullin: a peptide superfamily. 1997
- 126/78 09402770 97346111 PMID: 9202531
Same-day thyroid surgery: an analysis of safety, cost savings, and outcome. Jul 1997
- 126/79 09356270 97290598 PMID: 9145244
Effects of an inhibitor of cathepsin L on bone resorption in thyroparathyroidectomized and ovariectomized rats. May 1997
- 126/80 093533827 97310193 PMID: 9221312
Acute hyperparathyroidism associated with follicular carcinoma in the thyroid: possible role of juvenile cervical irradiation. Description of a case] Iperparatiroidismo acuto associato a carcinoma follicolare della tiroide: possibile ruolo dell'irradiazione cervicale giovanile. Descrizione di un caso clinico. Mar 1997
- 126/81 09339826 97324806 PMID: 9180911
Carcinoma of the thyroid in children: a 25-year experience. Aug 1997
- 126/82 09334258 97247907 PMID: 9094030
Heterotopic autotransplantation of parathyroid tissue in children undergoing total thyroidectomy. Mar 1997
- 126/83 09276587 97102604 PMID: 8943173
Familial hyperparathyroidism without multiple endocrine neoplasia. Jan 1997
- 126/84 09223203 96427228 PMID: 8678956
Prophylactic surgery for multiple endocrine neoplasia type IIa after genetic diagnosis: is parathyroid transplantation indicated? Sep 1996
- 126/85 09220766 96318976 PMID: 8734286
Detection of residual lesions and metastases of medullary thyroid cancer] Detection des lésions résiduelles et des métastases du cancer médullaire de la thyroïde. 1996
- 126/86 09144713 97063813 PMID: 8907680
Unusual types of thyroid neoplasms. Mar 1996
- 126/87 09115482 97064898 PMID: 8908448
Association of primary hyperparathyroidism and non-medullary thyroid cancer. Sep 1996
- 126/88 09030286 96301273 PMID: 8766940
Graves' disease in children and adolescents. Late results of surgical treatment. Jun 1996
- 126/89 09019074 96427227 PMID: 8678955
Primary hyperparathyroidism in multiple endocrine neoplasia type IIa: retrospective French multicentric study. Groupe d'Etude des Tumeurs a Calcitonine (GETC, French Calcitonin Tumors Study Group). French Association of Endocrine Surgeons. Sep 1996
- 126/90 09019072 96427225 PMID: 8678953
Surgical treatment of Graves' disease in patients younger than 18 years. Sep 1996
- 126/91 08952467 96316843 PMID: 8712911
Reoperation for persistent or recurrent primary hyperparathyroidism. Aug 1996
- 126/92 08933297 96228441 PMID: 8683653
Imaging and surgery in hyperparathyroidism: a 10-year review. Apr 1996
- 126/93 08930765 96288074 PMID: 8679414
[Total thyroidectomy with preservation of the parathyroid glands] La tiroïdectomie totale con conservazione delle paratiroidi. Jan-Feb 1996
- 126/94 08929221 96234536 PMID: 8677042
The choice of the intervention in the surgical treatment of nontoxic diffuse multinodular goiter] La scelta dell'intervento nel trattamento chirurgico del gozzo multinodulare diffuso non tossico. Jan-Feb 1996
- 126/95 08828776 96118989 PMID: 8581954
Morphological study of amelogenesis in the rat lower incisor after thyro-parathyroidectomy, parathyroidectomy and thyroidectomy. Jan 1996
- 126/96 08825191 96144649 PMID: 8560383
Progression of postoperative residual medullary thyroid carcinoma as monitored by plasma calcitonin levels. Jan 1996
- 126/97 08750274 96308582 PMID: 8712598
Surgical treatment of differentiated thyroid carcinomas: results in 440 cases operated between 1970 and 1994] Trattamento chirurgico dei carcinomi differenziali della tiroide: risultati su 440 casi operati dal 1970 al 1994. Nov-Dec 1995
- 126/98 08664445 96089017 PMID: 7491518
Laparoscopic detection of hepatic metastases in patients with residual or recurrent medullary thyroid cancer. Dec 1995
- 126/99 08658412 96079753 PMID: 7483076
Surgical treatment of goiter at a central hospital. A consecutive study with special emphasis on surgical complications] Kirurgisk behandling af struma på et centralsygehus. En konsekutiv opfølgelse specielt med henblik på kirurgiske komplikationer. Oct 23 1995